


EVALUATION OF THE RADIONUCLIDE CONCENTRATIONS IN SOIL AND PLANTS FROM THE 1975 TERRESTRIAL SURVEY OF BIKINI AND ENEU ISLANDS

C. S. Colsher, W. L. Robison, and P. H. Gudiksen

January 21, 1977

Prepared for U.S. Energy Research & Development
Administration under contract No. W-7405-Eng-48



5010073

NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Energy Research & Development Administration, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately-owned rights.

NOTICE

Reference to a company or product name does not imply approval or recommendation of the product by the University of California or the U.S. Energy Research & Development Administration to the exclusion of others that may be suitable.

Printed in the United States of America

Available from

National Technical Information Service

U S Department of Commerce

5285 Port Royal Road

Springfield, VA 22161

Price Printed Copy \$, Microfiche \$3.00

| Page Range | Domestic Price | Page Range | Domestic Price |
|------------|----------------|------------|----------------|
| 001-025 | \$ 3.50 | 326-350 | 10.00 |
| 026-050 | 4.00 | 351-375 | 10.50 |
| 051-075 | 4.50 | 376-400 | 10.75 |
| 076-100 | 5.00 | 401-425 | 11.00 |
| 101-125 | 5.50 | 426-450 | 11.75 |
| 126-150 | 6.00 | 451-475 | 12.00 |
| 151-175 | 6.75 | 476-500 | 12.50 |
| 176-200 | 7.50 | 501-525 | 12.75 |
| 201-225 | 7.75 | 526-550 | 13.00 |
| 226-250 | 8.00 | 551-575 | 13.50 |
| 251-275 | 9.00 | 576-600 | 13.75 |
| 276-300 | 9.25 | 601-up | * |
| 301-325 | 9.75 | | |

*Add \$2.50 for each additional 100 page increment from 601 to 1,000 pages,
add \$4.50 for each additional 100 page increment over 1,000 pages

5010014



LAWRENCE LIVERMORE LABORATORY

University of California Livermore, California 94550

UCRL-51879 Part 3

**EVALUATION OF THE RADIONUCLIDE CONCENTRATIONS
IN SOIL AND PLANTS FROM THE 1975 TERRESTRIAL
SURVEY OF BIKINI AND ENEU ISLANDS**

C. S. Colsher, W. L. Robison,
and P. H. Gudiksen

January 21, 1977

5010R75

BLANK PAGE

5010076

Contents

| | |
|---|-----|
| Abstract | 1 |
| Introduction | 2 |
| Methods | 4 |
| Environmental Sampling and Laboratory Analysis | 4 |
| Data Analysis | 5 |
| Results and Discussion | 12 |
| Surface Soil Survey | 12 |
| Soil Profiles | 14 |
| Prediction of Plant Uptake | 17 |
| Concentration Factors | 18 |
| Concentration Ratios | 28 |
| Summary and Conclusions | 31 |
| Acknowledgment | 36 |
| References | 38 |
| Appendix A. Soil Profiles of Bikini and Eneu Islands | A-1 |
| Appendix B. The Distribution of Radionuclides with Depth in Soil Profiles of Bikini and Eneu Islands | B-1 |
| Appendix C. Geographical Distribution of Radioactivity in the Surface Soil (0 to 15 cm) of Bikini and Eneu Islands | C-1 |
| Appendix D. Concentrations in Soil and Vegetation on Bikini and Eneu Islands . . . (microfiche in pocket on inside back cover) | |

BLANK PAGE

5016078

EVALUATION OF THE RADIONUCLIDE CONCENTRATIONS IN SOIL AND PLANTS FROM THE 1975 TERRESTRIAL SURVEY OF BIKINI AND ENEU ISLANDS

Abstract

In June 1975, personnel from LLL and from other laboratories and agencies conducted a radiological survey of the terrestrial environment of Bikini and Eneu Islands (Bikini Atoll) to evaluate the potential radiation dose to the returning Bikini population. In this report, we present measurements of the radionuclide concentration in soil profiles and in dominant species of edible and nonedible, indicator plants. We also describe the use of these data to derive relationships to predict the plant uptake of radionuclides from soil.

Approximately 620 soil and vegetation samples from Bikini and Eneu Islands were analyzed by Ge(Li) gamma spectrometry and by wet chemistry. The predominant radionuclides in these samples are ^{60}Co , ^{90}Sr , ^{137}Cs , $^{239,240}\text{Pu}$, ^{241}Pu , and ^{241}Am . In general, the radionuclide concentrations in soil from Eneu Island and from the four areas of Bikini Island appear to approximate log-normal distributions. The median surface-soil concentrations (pCi/g) of Eneu Island (0.067 for ^{60}Co , 4.1 for ^{90}Sr , 2.9 for ^{137}Cs , 0.25 for $^{239,240}\text{Pu}$, and 0.22 for ^{241}Am) are ten times lower than those measured on Bikini

Island (0.86 for ^{60}Co , 76 for ^{90}Sr , 43 for ^{137}Cs , 3.0 for $^{239,240}\text{Pu}$, and 2.4 for ^{241}Am). We found that radioactivity is unevenly distributed over the surface of these islands and that the distribution of activity with soil depth varies greatly in different parts of the islands. Concentrations in the soil of ^{90}Sr and ^{137}Cs are greater than concentrations of ^{241}Am and $^{239,240}\text{Pu}$ which, in turn, are greater than concentrations of ^{60}Co .

To quantitatively evaluate the plant uptake of ^{90}Sr , ^{137}Cs , and $^{239,240}\text{Pu}$, we develop soil-plant concentration factors as well as leaf-leaf and fruit-leaf concentration ratios for indicator and edible plant species from the same location. In general, the concentration factors for ^{137}Cs in terrestrial vegetation are greater than those for ^{90}Sr . The concentration factors for both of these nuclides exceed those for $^{239,240}\text{Pu}$ by one to two orders of magnitude (10 to 100 times). For ^{90}Sr and $^{239,240}\text{Pu}$, nuclide uptake by fruit is less than that by mature leaves; however, the opposite is true for ^{137}Cs . The relative contribution of the individual plant species to the internal dose to man

SOIL PAPER

varies with the nuclide under consideration. Thus, we also describe the use of concentration factors and

concentration ratios to predict nuclide concentrations in fruit from those observed in soil or leaves.

Introduction

Since the termination of nuclear testing on Bikini Atoll in 1958, periodic environmental surveys have been conducted to evaluate the radiological status of the atoll. The early surveys of Bikini Atoll as well as the recent survey and assessment at nearby Enewetak Atoll indicate that concentrations of radionuclides in certain terrestrial foods are relatively high, suggesting that the terrestrial foodchain could be a major exposure pathway.¹⁻⁴ In June 1975, Bikini Atoll was resurveyed to determine the residual radioactivity in the terrestrial environments of Bikini and Eneu Islands, the two main islands of the atoll (Fig. 1). The 1975 survey included measurement of environmental gamma-ray exposure rates and the collection and analysis of samples of soil, ground water, cistern water, and vegetation for use in assessing the internal dose via various ingestion pathways. (The dose from external gamma exposure and the radionuclide concentrations in cistern and ground water have been previously reported.^{5,6})

The longer-lived fission and activation products are the nuclides of primary concern at Bikini Atoll. Previous studies have shown that because of their long half lives and large inventories, ⁹⁰Sr, ¹³⁷Cs, and ^{239,240}Pu contribute nearly all the population dose from the terrestrial pathway.^{7,8} In this study, our major emphasis is on ⁹⁰Sr, ¹³⁷Cs, and ^{239,240}Pu. However, the results obtained for ⁶⁰Co and ²⁴¹Am are also included because ⁶⁰Co is widely distributed and is present in the marine pathway and because the concentration of ²⁴¹Am is still increasing slightly due to the decay of ²⁴¹Pu.

In this report, we describe the results of the soil and vegetation studies of the 1975 survey. We also discuss the use of the data to derive relationships that predict the plant uptake of nuclides from soil on Bikini Atoll. Geometric mean values of ⁶⁰Co, ⁹⁰Sr, ¹³⁷Cs, ^{239,240}Pu, and ²⁴¹Am surface-soil concentrations are developed for Eneu Island and for each of the four areas into which Bikini Island was divided. We also




Fig. 1. The Bikini Atoll.

analyzed soil profiles to investigate the distribution of activity with soil depth.

Soil-plant concentration factors and soil-plant regression equations, together with leaf-leaf and leaf-fruit concentration ratios, are calculated for ^{90}Sr , ^{137}Cs , and $^{239,240}\text{Pu}$ in edible and indicator plants as well as in soil from the associated sampling site. From our evaluation of these data, a method is developed for predicting the nuclide concentrations in edible plants at a

given location from the determined nuclide concentrations in soil at the same location. The predicted nuclide concentrations in edible plants subsequently serve as input data to predict the internal dose from ingested terrestrial foods. This dose assessment is presented in a separate report.⁹

At Bikini Atoll, the uptake by plants of radioactive material from the soil is the principal source of foodchain contamination. Thus, soil is both the convenient and the logical starting point for a prediction of

radionuclide concentration in terrestrial plants. Soil-plant concentration factors or soil-plant regression equations are commonly used for a quantitative comparison of the capacity of different plant species and various plant organs to accumulate radionuclides through soil-root uptake.¹⁰ Prediction of radionuclide uptake by edible plants is needed to convert

the measured environmental soil concentrations into the potential dose to man from the soil-plant pathway. Where fruit samples are not available, correlations between the concentrations in leaves and fruit of a particular plant species or between concentrations in leaves of indicator and edible plant species enable us to predict plant uptake from soil.

Methods

ENVIRONMENTAL SAMPLING AND LABORATORY ANALYSIS

The objective of the surface-soil survey was to define the distribution of radioactivity within the soil on Bikini and Eneu Islands. This survey was conducted in a manner similar to that used at Enewetak Atoll.¹ The number of soil samples collected on each island and within specific areas on the island was a function of the anticipated radioactivity levels, the various housing locations under consideration, and the expected living patterns of the future inhabitants. Thus, Eneu Island, because of its low and homogeneous activity levels was sampled less densely than Bikini Island. Bikini Island has elevated and more variable activities and was divided into four distinct areas, each of which could be used for future village sites. Sampling sites were

selected by superimposing a rectangular-grid network over an aerial photograph of each island and randomly choosing the grid squares to be sampled within each specific area of interest. The surface-sample locations for the islands are shown in Fig. 2. These samples were taken with a coring tool (a steel pipe, 30 cm² in cross-section) to a depth of 15 cm. The surrounding soil was scraped away and a cutting tool (a flat piece of steel) was inserted underneath the cover, freeing the sample.

In addition to the surface-soil sampling program, vegetation and associated soil profiles (soil profiles taken from the same location as the plant) were collected wherever suitable plant species were located on Bikini and Eneu Islands (see Fig. 3, Appendices A and B). Leaves,

fruit, roots, litter, and stems of edible (*Pandanus*, breadfruit, coconut, papaya, banana, and squash) and non-edible indicator (*Scaevola* and *Messerschmidia*) plants were collected when available. We attempted to take at least one soil profile, and preferably as many as three, through the root zone of each sampled plant. In addition to the soil profiles taken through the root zone of sampled vegetation, other soil profiles were collected on a random basis on both Bikini and Eneu Islands. The geographical locations of these profiles are also shown in Fig. 3.

All profile samples were taken from pits dug with a backhoe. After the pit was dug, the sidewalls were carefully cut back a few centimeters to ensure a clean, undisturbed profile. For each profile, 100-cm² samples were collected from the sidewall at increments of 5 cm in the upper part and of 10 to 20 cm through the lower part of the profile. Total depth for profile samples varied from 25 to 105 cm.


Vegetation and litter samples were carefully selected and classified by age. For example, collected leaves were classified as young, mature, and senescent. Two ages of litter were readily identifiable and were collected accordingly. However, fruit samples representing different stages of growth were unavailable.

All samples were handled separately, placed in plastic bags, and sent to LLL for processing and analysis. All soil and vegetation samples were analyzed both by Ge(Li) gamma spectroscopy and by wet chemistry for the following radionuclides (see Ref. 12): ^{40}K , ^{55}Fe , ^{60}Co , ^{90}Sr , ^{106}Ru , $^{102\text{m}}\text{Rh}$, $^{110\text{m}}\text{Ag}$, ^{125}Sb , ^{133}Ba , ^{137}Cs , ^{144}Ce , ^{152}Eu , ^{155}Eu , ^{207}Bi , ^{228}Ra , ^{235}U , ^{238}Pu , ^{239}Pu , ^{240}Pu , ^{241}Pu , and ^{241}Am .

DATA ANALYSIS

The surface-soil (0 to 15 cm) activities appear to approximate log-normal distributions and thus, we calculated geometric means of ^{60}Co , ^{90}Sr , ^{137}Cs , $^{239,240}\text{Pu}$, and ^{241}Am concentrations in soil for Eneu Island as a whole and for the four areas of interest on Bikini Island. For each profile collected, we plotted (on semilog paper) the concentrations of the selected nuclides as a function of depth. The profile data were compared in an attempt to characterize the different areas of the islands (see Appendices A and B).

Because they are the major contributors to the dose from ingestion of terrestrial foods (Refs. 1, 7, 8), ^{90}Sr , ^{137}Cs , and $^{239,240}\text{Pu}$ were selected for more detailed analysis. We calculated concentration factors for these nuclides from measured






Fig. 2. Sampling sites of the Bikini (a) and Eneu (b) surface-soil sampling program.




Fig. 3. Sampling sites of the Bikini (a) and Eneu (b) soil profile and vegetation samples.

concentrations in plant samples and from the average measured concentration in the associated 0- to 25-cm soil profile. We define the concentration factor, CF, as

$$CF = \frac{\text{pCi/g dry plant}}{\text{pCi/g dry soil}}$$

To reduce the variability in average soil concentrations (used to calculate the concentration factor), we used a 0- to 25-cm soil profile that encompasses a large fraction of the effective absorptive root zone rather than the deeper 0- to 55-cm profile that encompasses the entire root zone. Concentration factors calculated on the basis of the average soil concentration in the upper 25 cm of the profile are somewhat greater but do not differ substantially from those based on the deeper profiles (Tables 1 and 2). All concentration factors reported here are therefore those derived from average 0- to 25-cm soil concentrations.

The average 0- to 25-cm soil concentration is calculated as the weighted geometric mean for the separate concentrations, measured at various increments throughout the profile. Concentration values less than the minimum detection limit are set equal to the detection limit, following the U.S. Environmental Protection Agency technique.

Whenever the number of samples is large enough, soil concentrations are plotted against concentrations in plants from the same sampling site; the results are analyzed with linear regression methods. These linear regression results are "statistically significant" at the 0.1-level of a standard F test. For each combination of nuclide, plant organ, and species considered, statistically significant regression equations are compared to the median of the calculated concentration factors and a single representative concentration factor is assigned.

Predictions of radionuclide levels in foodstuffs can be made from concentration factors if measured soil concentrations are available; however, concentration ratios are also needed if the only available data are from mature leaf samples. The concentration ratio is defined as the ratio of the concentration in fruit to the concentration in leaves of the same species; or, as the concentration in leaves of one species to the concentration in leaves of another species. We calculated preliminary ratios for all available species from the 1975 Bikini survey. However, because of the small number of samples involved, a statistical analysis of these results was not possible.

Table 1. Average radionuclide concentration for 0- to 25-cm and deeper soil profiles.

| Location ^a | Average Soil Concentration, pCi/g dry weight | | | |
|---------------------------|--|-------------------|-------------------|-------------------|
| | ⁹⁰ Sr | ¹³⁷ Cs | ²³⁹ Pu | ²⁴⁰ Pu |
| T0001 (0-25) ^b | 81 | 45 | 4.3 | 4.7 |
| (0-40) | 70 | 27 | 1.6 | 1.8 |
| T0051 (0-25) | 202 | 150 | 9.2 | 10 |
| (0-45) | 208 | 165 | 9.2 | 10 |
| T0061 (0-25) | 42 | 28 | 1.4 | 1.6 |
| (0-55) | 150 | 85 | 5.1 | 5.7 |
| T0062 (0-25) | 67 | 31 | 2.3 | 2.5 |
| (0-45) | 80 | 37 | 1.3 | 1.5 |
| T0081 (0-25) | 126 | 43 | 2.2 | 2.5 |
| (0-45) | 70 | 15 | 0.53 | 0.64 |
| T0121 (0-25) | 89 | 50 | 2.0 | 2.2 |
| (0-55) | 34 | 25 | 0.52 | 0.60 |
| T0161 (0-25) | 27 | 34 | 1.5 | 1.7 |
| (0-55) | 30 | 25 | 1.1 | 1.2 |
| T0181 (0-25) | 94 | 13 | 2.4 | 2.8 |
| (0-55) | 62 | 5 | 0.56 | 0.70 |
| T0191 (0-25) | 36 | 23 | 0.99 | 1.2 |
| (0-55) | 28 | 18 | 0.87 | 1.0 |
| T0241 (0-25) | 3.7 | 3.3 | 0.42 | 0.46 |
| (0-45) | 4.7 | 2.8 | 0.49 | 0.47 |
| T0251 (0-25) | 7.3 | 7.9 | 0.24 | 0.27 |
| (0-45) | 9.8 | 5.7 | 0.42 | 0.45 |
| T0261 (0-25) | 7.8 | 4.6 | 0.41 | 0.47 |
| (0-45) | 6.8 | 1.8 | 0.23 | 0.38 |
| T0271 (0-25) | 1.0 | 0.88 | 0.12 | 0.14 |
| (0-45) | 0.62 | 0.40 | --- | --- |
| T0301 (0-25) | 16 | 10 | 1.1 | 1.2 |
| (0045) | 20 | 12 | 1.6 | 1.5 |

^a Sample locations are shown in Fig. 2.

^b Depth of soil profile in centimeters.

^c Not detected.

Table 2. Soil-mature leaf concentration factors calculated for 0- to 25-cm and deeper soil profiles.

| Location ^a | Concentration Factor, (pCi/g dry leaf)/(pCi/g dry soil) | | | |
|-----------------------|---|-------------------|-------------------|-------------------|
| | ⁹⁰ Sr | ¹³⁷ Cs | ²³⁹ Pu | ²⁴⁰ Pu |
| T0001 (0-25) | 1.0 | 17 | 0.01 | 0.01 |
| (0-40) | 1.2 | 29 | 0.038 | 0.01 |
| T0051 (0-25) | 0.94 | 0.30 | 0.050 | 0.0053 |
| (0-40) | 0.92 | 0.27 | 0.050 | 0.0052 |
| T0061 (0-25) | 5.3 | 2.5 | 0.043 | 0.045 |
| (0-55) | 0.15 | 0.82 | 0.012 | 0.013 |
| T0062 (0-25) | 2.8 | 2.2 | 0.049 | 0.045 |
| (0-55) | 3.3 | 1.9 | 0.028 | 0.013 |
| T0081 (0-25) | 1.4 | 0.79 | 0.019 | 0.0494 |
| (0-45) | 2.4 | 2.3 | 0.078 | 0.029 |
| T0121 (0-25) | 1.2 | 2.9 | 0.030 | 0.021 |
| (0-55) | 2.5 | 5.9 | 0.066 | 0.084 |
| T0161 (0-25) | 0.22 | 13 | 0.018 | 0.014 |
| (0-55) | 0.19 | 18 | 0.024 | 0.054 |
| T0181 (0-25) | 0.11 | 35 | 0.012 | 0.018 |
| (0-55) | 0.17 | 9.3 | 0.051 | 0.0245 |
| T0191 (0-25) | 0.56 | 17 | 0.025 | 0.013 |
| (litter) (0-55) | 0.72 | 23 | 0.029 | 0.051 |
| T0251 (0-25) | 0.30 | 1.1 | --- | --- |
| (0-45) | 0.19 | 1.3 | --- | --- |
| T0261 (0-25) | 0.16 | 3.9 | 0.010 | 0.013 |
| (0-45) | 0.12 | 5.4 | 0.0059 | 0.069 |
| T0261 (0-25) | 0.099 | 2.6 | --- | --- |
| (0-45) | 0.20 | 6.6 | --- | --- |
| T0271 (0-25) | --- | 1.6 | --- | --- |
| (0-45) | --- | 36 | --- | --- |
| T0301 (0-25) | 0.11 | 2.3 | --- | --- |
| (0-45) | 0.085 | 2.0 | --- | --- |

^a Sample locations are shown in Fig. 2.

^b No data.

Results and Discussion

SURFACE SOIL SURVEY

Although more samples are available from the 1975 Bikini survey than from any previous survey (Table 3), there is little consistency in the geographical distribution of ^{60}Co , ^{90}Sr , ^{137}Cs , $^{239,240}\text{Pu}$, and ^{241}Am on Bikini and Eneu Islands (see Appendix C). The maps and overlays in Appendix C present the activities of these radionuclides in picocuries per gram of dry soil over the sites from which the samples were collected. A list of concentrations of all detectable nuclides for each sampling site is given in Appendix D (microfiche included in pocket on inside back cover). A dry-soil density of 1.5 g/cm³ may be used to convert the integrated profile data into activity per unit area. However, some caution must be exercised in such calculations because a significant fraction of the total activity may be located below the sampling depth.

Table 4 presents the means of the surface-soil concentrations of the dominant nuclides for Eneu Island and for the four areas of interest on Bikini Island. The values for Eneu are consistently ten times lower than concentrations for any part of Bikini Island. As expected, on Bikini Island,

Area 1 shows the lowest soil concentration, since it is an exposed beach area that has been cleared for housing. Data from Bikini⁸ and Enewetak¹² have revealed that soil activity is directly related to the amount of vegetation present in the area surrounding the sampling site. One possible reason for this is that a heavy vegetative cover can protect the underlying soil, minimizing the effects of weathering processes (e.g., wind and rain erosion) that transport surface activity through the soil column to the water lens. Follow-up field work at Enewetak Atoll has also shown that, in heavily vegetated areas, litter increases the soil retention of radionuclides.¹³

Although soil concentrations of radionuclides in Area 2 appear to be higher for ^{90}Sr and $^{239,240}\text{Pu}$ than in any other area on Bikini, statistical analysis of the ^{90}Sr concentrations for each of the four areas on Bikini, using the Mann-Whitney nonparametric test, shows no significant difference between the concentrations in the various areas. However, a more extensive analysis is needed to better define the real differences in concentrations in the various areas.

Table 3. Number of vegetation and soil samples in various Bikini surveys.

| Species, Organ | Number of Samples | | | | | |
|--|------------------------------|-----------------------------|-------------------|-----------------|--------------------------|----------------|
| | ^{90}Sr | | ^{137}Cs | | $^{239},^{240}\text{Pu}$ | |
| | 64-74 ^a Survey | 1975 ^b Survey | 64-74 Survey | 1975 Survey | 64-74 Survey | 1975 Survey |
| <i>Pandanus</i> , leaves ^c fruit | 6 | 6 (6) ^d 1 (1) | 4 | 6 (6) 1 (1) | 5 (5) 1 (1) | |
| Papaya, leaves fruit | | 8 (4) 4 (4) | | 8 (4) 4 (4) | 8 (4) 4 (4) | |
| Breadfruit, leaves fruit | 0 | 2 (2) 1 (1) | 0 | 2 (2) 1 (1) | 2 (2) 1 (1) | |
| Banana, leaves | | 3 (3) | | 3 (3) | 3 (3) | |
| Coconut, leaves fruit | 5 | 22 (8) 6 (6) | 48 | 22 (8) 6 (6) | 22 (8) 6 (6) | |
| <i>Scaevola</i> , leaves fruiting body | | 8 (2) 1 (1) | | 8 (2) 1 (1) | 8 (2) 1 (1) | |
| <i>Messerschmidia</i> , leaves | | 6 (3) | | 6 (3) | 6 (3) | |
| Soil Profiles | 5 | 42 | 22 | 42 | 42 | |
| Soil, top 15 cm | 21 | 196 | 176 | 196 | 196 | |

^a Data from 1974 Bikini draft (unpublished).^b Data from this report.^c Leaves include both mature and young specimens.^d Number of samples that have directly associated soil profiles.

Table 4. Median surface soil concentrations (0 to 15 cm).

| Nuclide | Soil Concentration, pCi/g dry weight | | | | | Bikini Island ^a | Eneu Island |
|------------------------|--------------------------------------|----------|-----------|-----------|------------|----------------------------|-------------|
| | Area 1 | Area 2 | Area 3 | Area 4 | | | |
| ⁶⁰ Co | 0.59 (51) ^b | 0.98 (6) | 0.94 (32) | 0.92 (87) | 0.86 (176) | 0.067 (66) | |
| ⁹⁰ Sr | 41 (35) | 126 (5) | 69 (18) | 68 (70) | 76 (128) | 4.1 (73) | |
| ¹³⁷ Cs | 34 (51) | 43 (6) | 48 (33) | 48 (88) | 43 (178) | 2.9 (68) | |
| ^{239, 240} Pu | 2.3 (70) | 4.5 (10) | 2.1 (34) | 3.0 (140) | 3.0 (254) | 0.25 (146) | |
| ²⁴¹ Am | 1.8 (51) | 3.7 (6) | 2.7 (31) | 2.7 (87) | 2.4 (175) | 0.22 (68) | |

^a Arithmetic average of soil concentration in Areas 1-4 on Bikini Island.

^b Number of samples taken.

SOIL PROFILES

Soil profiles from different parts of Bikini and Eneu Islands show a wide range of activity distributions with depth. (A listing of the concentrations measured for each profile is given in Appendix A and the data are presented graphically in Appendix B.) As noted by Held,¹⁴ different plant-soil environments exhibit different vertical patterns of nuclide migration. The nearly complete disruption of the upper soil layers at Bikini Atoll by clearing, construction, and testing over the past 30 years as well as by agricultural practices initiated more recently has created a variety of plant-soil environments. Thus, the inhomogeneity of the soil on these islands is not surprising. However,

because of this inhomogeneity, generalizations are not very meaningful, and these islands cannot be characterized by "average vertical profiles" with which to formulate cleanup criteria and to estimate dose.

The four basic types of profiles delineated at Enewetak Atoll¹² are all present on Bikini Island (see Fig. 4). Although we could not identify any particular profile type for extensive areas on Bikini Island, specific locations can be assigned "typical" profiles for predictive purposes. For example, on Bikini Island a group of samples taken in close proximity to one another (T0091, T0101, T0111, T0121, and T0131) show generally decreasing activity levels with depth despite




Fig. 4. Basic profile types from Gudiksen¹¹: (a) erratic, (b) sharp drop-off, (c) uniform throughout, and (d) increasing then decreasing.

some variability in pattern in the upper parts of the profiles (see Figs. 5-9; Appendices A and B). Other locations, limited in area, can be assigned different "typical" profiles.

In general, profiles from Bikini Island show decreasing activity levels with depth. In contrast, those from Eneu Island exhibit a pattern of uniform or slowly decreasing activity levels from surface to total-sampled depth. The variations seen in the profiles on Bikini may be the result

of the location of organic layers in the profile. Because organic matter tends to concentrate radioactivity,¹⁵ nonuniform patterns of radionuclide concentration may result from organic layers that have been buried recently by construction and rehabilitation activities.

Although it is difficult to generalize about patterns of activity distribution, the relative concentrations of the dominant nuclides show a consistent trend: ⁹⁰Sr and ¹³⁷Cs >




Fig. 5. Correlation of the ⁹⁰Sr concentration in mature coconut leaves with the concentration of ⁹⁰Sr in the soil at the same site.




Fig. 6. Correlation of the ^{137}Cs concentration in mature coconut leaves with the concentration of ^{137}Cs in the soil at the same site.

$^{239,240}\text{Pu}$ and $^{241}\text{Am} > ^{60}\text{Co}$. The concentration of ^{90}Sr on Bikini Island is usually twice that of ^{137}Cs , ten times that of $^{239,240}\text{Pu}$ and ^{241}Am , and thirty to forty times that of ^{60}Co (Table 4). As mentioned previously, soil concentrations on Eneu are about ten times lower than those on Bikini Island for all the radionuclides considered.

PREDICTION OF PLANT UPTAKE

Average concentrations in the 0- to 25-cm portions of the soil profiles are combined with measured concentrations in plants to predict the uptake of various radionuclides from the soil. In general, these plant-soil relationships from the 1975 Bikini survey confirm the results of previous surveys; however, the relationships are often statistically insignificant.




Fig. 7. Correlation of the $^{239,240}\text{Pu}$ concentration in mature coconut leaves with the concentration of $^{239,240}\text{Pu}$ in the soil at the same site.

In surveys where the number of samples considered for any one case is small, relationships that appear to be statistically insignificant are often extremely significant, since sampling errors may dominate the explanatory variables. Although statistical analysis of a larger number of samples is necessary to

verify the results, we recommend the use of the general plant-soil relationships developed in this study for subsequent dose assessments.

Concentration Factors

Soil profiles with uniform patterns of nuclide migration are seldom found at Bikini Atoll as a




Fig. 8. Correlation of the ^{90}Sr concentration in mature *Scaevola* and *Messerschmidia* leaves with the concentration of ^{90}Sr in the soil at the same site.

result of the complete disruption of the upper soil layers by clearing, construction, and testing over the past 30 years as well as by more recent agricultural practices. To determine the soil concentrations of nuclides that are actually available to the root system of a specific plant, we sampled soil profiles in direct contact with the root system. The two replicate samples of soil profiles show minimal variation,

regardless of the side of the plant from which they were taken (Table 5). In contrast, profiles in the general area but not in direct contact with the root system of the plant sample are highly variable (Table 6).

Tables 7 and 8 present the range and median values of concentration factors calculated for vegetation and soil sampled from the same location. Table 9 compares the information from these tables with the same information




Fig. 9. Correlation of the ^{137}Cs concentration in mature *Scaevola* and *Messerschmidia* leaves with the concentration of ^{137}Cs in the soil at the same site.

for concentration factors calculated for mature *Scaevola* and coconut leaf samples for which no soil samples from the same location are available. We selected mature *Scaevola* and coconut leaves for this comparison because they provide the largest number of samples in both the associated and unassociated categories. A comparison of the ranges in Table 9 shows the importance of using associated plant-soil data (data from the same sampling

site). Concentration factors calculated from unassociated plant and soil factors show a variation of three orders of magnitude in the case of ^{137}Cs uptake by mature *Scaevola* leaves while concentration factors calculated from associated data vary by one order of magnitude or less. These results agree with the wide range of concentration factors calculated in previous surveys from unassociated plant and soil samples.^{1,4}

Table 5. Radionuclide concentrations in replicate 0- to 25-cm soil profiles.

| Location ^a | Average Soil Concentration, pCi/g dry weight | | | |
|-----------------------|--|-------------------|-------------------|-------------------|
| | ⁹⁰ Sr | ¹³⁷ Cs | ²³⁹ Pu | ²⁴⁰ Pu |
| <u>Group 1</u> | | | | |
| T0001 | 81 | 45 | 4.3 | 4.7 |
| T0002 | 92 | 29 | 4.6 | 5.1 |
| T0003 | 101 | 44 | 5.0 | 5.5 |
| <u>Group 2</u> | | | | |
| T0061 | 42 | 28 | 1.4 | 1.6 |
| T0062 | 68 | 31 | 2.6 | 2.5 |

^aSampling sites are shown in Fig. 2.

Table 6. Radionuclide concentrations in 0- to 25-cm soil profiles taken from the same general area.

| Location ^a | Average Soil Concentration, pCi/g dry weight | | | |
|-----------------------|--|-------------------|-------------------|-------------------|
| | ⁹⁰ Sr | ¹³⁷ Cs | ²³⁹ Pu | ²⁴⁰ Pu |
| <u>Group 1</u> | | | | |
| T0091 | 81 | 55 | 2.5 | 2.8 |
| T0101 | 32 | 35 | 0.27 | 0.29 |
| T0111 | 81 | 87 | 3.3 | 3.7 |
| T0121 | 89 | 50 | 2.0 | 2.2 |
| T0131 | 86 | 13 | 0.53 | 0.64 |
| <u>Group 2</u> | | | | |
| T0031 | 6.8 | 8.6 | 0.18 | 0.20 |
| T0041 | 35 | 26 | 0.58 | 0.69 |
| T0051 | 202 | 150 | 9.2 | 10 |
| T0061 | 42 | 28 | 1.2 | 1.6 |
| T0071 | 127 | 86 | 4.6 | 5.1 |

^a Sampling sites are shown in Fig. 2.

Table 7. Soil-to-plant uptake of ^{90}Sr and ^{137}Cs from plants and soils sampled at the same location.

| Species, Organ | No. of Samples | Concentration Factor, $(\text{pCi/g dry plant}) / (\text{pCi/g dry soil})$ | | | No. of Samples | ^{137}Cs | | |
|--|----------------|--|---------|---------|----------------|-------------------|---------|--------|
| | | ^{90}Sr | Minimum | Maximum | | Minimum | Maximum | Median |
| <i>Scaevola</i> , mature leaves | 2 | 0.24 | 0.41 | 0.33 | 2 | 1.3 | 14 | 7.4 |
| <i>Messerschmidia</i> , mature leaves | 3 | 0.48 | 0.86 | 0.52 | 3 | 2.1 | 50 | 3.7 |
| Pooled <i>Scaevola</i> & <i>Messerschmidia</i> , mature leaves | 5 | 0.24 | 0.86 | 0.48 | 5 | 1.3 | 50 | 3.7 |
| Coconut, mature leaves | 7 | 0.099 | 0.38 | 0.16 | 8 | 1.1 | 16 | 3.0 |
| Coconut, "fruit" | 2 | 0.024 | < 0.018 | --- | 2 | 1.4 | 3.6 | 2.5 |
| Coconut, meat | 2 | < 0.019 | 0.026 | --- | 2 | 7.3 | 9.8 | 8.6 |
| Coconut, milk ^a | 2 | < 0.0084 | < 0.012 | --- | 2 | 0.90 | 1.4 | |
| <i>Pandanus</i> , mature leaves | 5 | 0.71 | 2.4 | 0.91 | 5 | 2.9 | 25 | 15 |
| <i>Pandanus</i> , green fruit | 1 | --- | --- | 0.53 | 1 | --- | --- | 0.054 |
| Papaya, mature leaves | 4 | 0.62 | 4.0 | 1.3 | 4 | 0.30 | 5.9 | 3.1 |
| Papaya, fruit | 4 | 0.12 | 0.85 | 0.43 | 4 | 1.9 | 18 | 3.2 |
| Breadfruit, mature leaves | 2 | 1.4 | 2.3 | 1.8 | 2 | 0.79 | 2.4 | 1.6 |
| Breadfruit, mature fruit | 1 | --- | --- | 0.76 | 1 | --- | --- | 7.0 |
| Banana, mature leaves | 2 | 0.48 | 1.1 | 0.73 | 2 | 0.33 | 0.54 | 0.42 |
| Squash, whole plant | 1 | --- | --- | 3.4 | 1 | --- | --- | 26 |
| Squash, seeds | 1 | --- | --- | 0.15 | 1 | --- | --- | 56 |

^a Coconut milk was measured and reported in pCi/ml wet weight which, for calculation of the concentration factor, was assumed to equal pCi/g wet weight. Thus, the concentration factor for coconut milk is in $(\text{pCi/g wet weight}) / (\text{pCi/g dry soil})$.

^b No data.

Table 8. Soil-to-plant uptake of $^{239},^{240}\text{Pu}$ from plants and soils sampled at the same location.

| Species, Organ | Concentration Factor, (pCi/g dry plant)/(pCi/g dry soil) | | | | | | | | | |
|--|--|-------------------|----------------|---------|---------|--------|----------------|---------|---------|--------|
| | ^{239}Pu | ^{240}Pu | No. of Samples | Minimum | Maximum | Median | No. of Samples | Minimum | Maximum | Median |
| <i>Scaevola</i> , mature leaves | 1 | --- | --- | --- | 0.0047 | 0.0047 | 1 | --- | --- | 0.0051 |
| <i>Messerschmidia</i> , mature leaves | 2 | 0.024 | 0.11 | 0.067 | 0.067 | 2 | 0.045 | 0.12 | 0.081 | |
| Pooled <i>Scaevola</i> & <i>Messerschmidia</i> , mature leaves | 3 | 0.0047 | 0.11 | 0.024 | 0.024 | 3 | 0.0051 | 0.12 | 0.045 | |
| Coconut, mature leaves | 4 | 0.010 | 0.022 | 0.015 | 0.015 | 4 | 0.0113 | 0.021 | 0.015 | |
| <i>Pandanus</i> , mature leaves | 4 | 0.0044 | 0.030 | 0.016 | 0.016 | 4 | 0.0043 | 0.015 | 0.014 | |
| Papaya, mature leaves | 4 | 0.0013 | 0.037 | 0.037 | 0.037 | 4 | 0.0053 | 0.041 | 0.026 | |
| Papaya, fruit | 2 | 0.0013 | 0.0021 | 0.0017 | 0.0017 | 2 | 0.0013 | 0.0023 | 0.0018 | |
| Breadfruit, mature leaves | 2 | 0.0063 | 0.019 | 0.013 | 0.013 | 2 | 0.0213 | 0.062 | 0.042 | |
| Banana, mature leaves | 2 | 0.0017 | 0.0054 | 0.0036 | 0.0036 | 2 | 0.0018 | 0.0066 | 0.0042 | |

^a No data

Table 9. Soil-mature leaf concentration factors calculated from associated^a and unassociated^b data.

| Nuclide, Species | Concentration Factor, (pCi/g dry plant)/(pCi/g dry soil) | | | | | | | |
|-------------------------------------|--|------------|---------|--------|----------------|--------------|---------|--------|
| | No. of Samples | Associated | | | No. of Samples | Unassociated | | |
| | | Minimum | Maximum | Median | | Minimum | Maximum | Median |
| ^{90}Sr , <i>Scaevola</i> | 2 | 0.24 | 0.41 | 0.33 | 4 | 0.048 | 4.3 | 1.8 |
| ^{90}Sr , coconut | 7 | 0.099 | 0.38 | 0.16 | 15 | 0.041 | 0.74 | 0.29 |
| ^{137}Cs , <i>Scaevola</i> | 2 | 1.3 | 14 | 7.5 | 4 | 0.073 | 39 | 7.7 |
| ^{137}Cs , coconut | 8 | 1.1 | 16 | 3.0 | 15 | 0.53 | 18 | 2.6 |
| ^{239}Pu , coconut | 4 | 0.011 | 0.022 | 0.015 | 12 | 0.0036 | 0.14 | 0.016 |
| ^{240}Pu , coconut | 4 | 0.011 | 0.021 | 0.015 | 12 | 0.0021 | 0.15 | 0.016 |

^a Plant and soil data sampled from the same site.

^b Plant and soil data sampled from different sites in the same general area.

Because the range of concentration factors in only two combinations of nuclide, plant part, and species from the associated soil-plant data (pooled *Scaevola*-*Messerschmidia* leaves for ^{137}Cs and papaya leaves for ^{239}Pu) varied by more than a factor of 20 (Table 10), we use the median concentration factors derived from the associated data in our predictive model.

Several reasons explain the variation of concentration factors calculated from associated plant and soil data, including differences in the physiochemical properties of the radionuclides under consideration, in soil type and chemical characteristics, in soil management practices, in irrigation practices, and in the physiology, age, and prior history of the plants sampled. It is impossible

Table 10. Maximum-to-minimum ratios of associated soil-plant concentration factors.

| Species, Organ | ^{90}Sr | Maximum-to-Minimum Ratio | | |
|--|------------------|--------------------------|-------------------|-------------------|
| | | ^{137}Cs | ^{239}Pu | ^{240}Pu |
| <i>Scaevola</i> , mature leaves | 1.7 | 10 | --- | ^a --- |
| <i>Messerschmidia</i> , mature leaves | 1.8 | 23 | 4.9 | 2.6 |
| Pooled <i>Scaevola</i> and <i>Messerschmidia</i> , mature leaves | 3.6 | 39 | 23 | 2.0 |
| <i>Pandanus</i> , mature leaves | 3.3 | 8.4 | 6.9 | 3.6 |
| Coconut, mature leaves | 3.8 | 14 | 2.1 | 1.8 |
| Coconut, "fruit" ^b | < 7.2 | 2.6 | --- | --- |
| Papaya, mature leaves | 6.5 | 20 | 30 | 7.9 |
| Papaya, fruit | 7.4 | 9.5 | 1.7 | 1.8 |
| Banana, mature leaves | 2.3 | 1.6 | 3.2 | 3.8 |
| Breadfruit, mature leaves | 1.7 | 3.0 | 3.0 | 2.9 |

^a Not detected.

^b "Fruit" includes both meat and milk.

to identify the specific cause of each variation but the variation can be reduced by carefully controlling sampling techniques and by increasing the number of samples.

Where fruit data are unavailable, concentration factors calculated from mature leaf data are used as the basis for predicting concentrations of radionuclides in food available to the returning Bikini population. Mature leaf concentration factors in conjunction with correlations between various species and between leaves and fruit of the same species enable us to predict concentrations in fruit from measured concentrations in leaves of indicator or edible plants.

We only report concentration factors for ^{90}Sr , ^{137}Cs , and $^{239,240}\text{Pu}$. As predicted from previous studies, the most effectively transferred radionuclide in the terrestrial environment is ^{137}Cs , although ^{90}Sr is often present in larger quantities in the soil of the atoll. This is partly explained by the differential solubilities of ^{90}Sr and ^{137}Cs in soil. Strontium-90 appears to be tied up as insoluble carbonates in the atoll soil and is thus less available to the plant. Cesium-137 is more soluble in the nonclay atoll soil; thus ^{137}Cs is more easily leached through the soil. Although ^{137}Cs is leached through the soil at

a faster rate than ^{90}Sr , ^{137}Cs is also readily absorbed and accumulated in organified soil horizons where there is a proliferation of plant roots and litter. This accumulation of ^{137}Cs in organified soil horizons renders it more available than ^{90}Sr for uptake in plants.

The concentration factors discussed in the subsequent paragraphs generally reflect this relationship between the uptake of ^{90}Sr and ^{137}Cs . For *Pandanus*, coconut, *Scaevola*, and *Messerschmidia* leaves and for the fruit of all species, low concentration factors are observed for ^{90}Sr as compared to ^{137}Cs . However, leaves of papaya, banana, and breadfruit show concentration factors for ^{90}Sr as high or higher than those for ^{137}Cs . Concentration factors for $^{239,240}\text{Pu}$ are generally 10 to 100 times lower than those for either ^{90}Sr or ^{137}Cs . Although they are often measured in soil, ^{60}Co , ^{241}Pu , and ^{241}Am are only occasionally detected in vegetation. For this reason, we did not calculate concentration factors for these three nuclides.

In the following sections, we discuss the specific concentration factors assigned to each species. Concentration factors are assigned solely on the basis of the median calculated concentration factors except for coconut and for pooled *Scaevola* and *Messerschmidia* leaves.

For these last two cases, we had enough samples to justify analysis by linear regression methods. Thus, the regression results are compared with the median concentration factor and a representative value is chosen for our models. The relationships between the relative uptake of different species are considered in a separate section on concentration ratios.

Coconut. Coconut is the most abundant species on Bikini and Eneu Islands and thus it was sampled more extensively than any other plant in the 1975 survey. Unfortunately, few coconut trees were bearing fruit so the bulk of these samples are leaves. Regression analysis comparing mature coconut leaves and soil sampled from the same location shows correlations that are significant at the 0.1 to 0.05 level for ^{90}Sr , at the 0.005 level for ^{137}Cs , and at the 0.1 level for $^{239,240}\text{Pu}$ (Figs. 5-7). Combining the results of this regression analysis with the median calculated concentration factors (Table II), we obtain final concentration factors for mature coconut leaves of 0.16 for ^{90}Sr , 3.0 for ^{137}Cs , and 0.015 for $^{239,240}\text{Pu}$.

Concentrations in both coconut milk and coconut meat were analyzed for the two samples from Bikini. When compared on a wet/wet or a dry/dry basis, there are no definitive patterns in the radionuclide concentrations of meat and milk taken from the

same location.* Because the 1972 Enewetak data show no consistent differences in the uptake of ^{90}Sr and ^{137}Cs by coconut milk and meat, the ^{90}Sr and ^{137}Cs concentrations in fresh coconut meat and fresh coconut milk are assumed to be equal. The concentration factors from the Bikini data (Table II) are within the range of those from the Enewetak survey,¹ so until more conclusive data are available, we have assigned a conservative concentration factor of 0.024 for ^{90}Sr and 2.5 for ^{137}Cs to both coconut meat and milk.

Pandanus -- Although the number of samples of mature *Pandanus* leaves is insufficient for statistical analysis, the concentration factors calculated from associated leaf-soil data for Bikini are within the range of those from Enewetak.¹ The median concentration factors of 0.91 for ^{90}Sr and 15.2 for ^{137}Cs are assigned to mature *Pandanus* leaves. The concentration factor calculated for the one green *Pandanus* fruit available from the Bikini survey (0.50 for ^{90}Sr and 5.4 for ^{137}Cs) is comparable to values from previous surveys and therefore, we

*Following the example of the Enewetak survey,¹ results are given in wet weight for coconut milk and in dry weight for coconut meat; coconut milk is assumed to be 95% water and coconut meat is assumed to be 50% water.

Table 11. Comparison of soil-root and soil-leaf concentration factors.

| Species | Concentration Factor, (pCi/g dry plant)/(pCi/g dry soil) | | | | | | |
|-----------------------|--|--------|-------------------|--------|-----------------------|--------|--|
| | ^{90}Sr | | ^{137}Cs | | $^{239,240}\text{Pu}$ | | |
| | Roots | Leaves | Roots | Leaves | Roots | Leaves | |
| <i>Pandanus</i> | 0.88 | 1.0 | 23 | 17 | 0.44 | 0.015 | |
| | 0.77 | 1.2 | 26.1 | 2.9 | 0.45 | 0.022 | |
| <i>Messerschmidia</i> | 0.59 | 0.48 | 33 | 50 | 0.44 | 0.35 | |
| Coconut | 0.16 | 0.11 | 3.4 | 2.3 | 0.085 | --- | |
| | 0.89 | --- | 16 | 0.88 | 0.11 | --- | |
| | 0.30 | 0.30 | 5.7 | 1.1 | 0.081 | --- | |
| | 0.89 | --- | 5.0 | --- | 0.38 | --- | |
| Banana | 0.42 ^a | 1.10 | 0.026 | 0.54 | 0.027 | 0.006 | |

^a Root and crown.

^b Below minimum detection limit.

consider it to be a valid sample and have used it in our dose predictions.

Breadfruit — Breadfruit were not available on Enewetak Atoll. Thus, data from the two samples collected during the 1975 Bikini survey provide the first directly measured concentration factors. As expected from the 1972 Bikini survey¹⁶ and from the stable element analysis of potassium and calcium in the Enewetak survey,¹ uptake of ^{90}Sr and ^{137}Cs by breadfruit is high and comparable to uptake by *Pandanus*. Preliminary concentration factors of 1.8 for ^{90}Sr , 1.6 for ^{137}Cs , and 0.027 for $^{239,240}\text{Pu}$ are assigned to mature breadfruit leaves. The concentration factors for the one mature

fruit sampled from this species are 0.76 for ^{90}Sr and 7.0 for ^{137}Cs .

Papaya — The ratio of maximum-to-minimum concentration factors calculated for mature papaya leaves is small; median concentration factors are 1.3 for ^{90}Sr , 3.1 for ^{137}Cs , and 0.0018 for $^{239,240}\text{Pu}$. Concentration factors of 0.43 for ^{90}Sr , 8.2 for ^{137}Cs , and 0.0018 for $^{239,240}\text{Pu}$ are calculated for the four papaya fruit available.

Banana — The only data from the 1975 Bikini survey on radionuclide concentrations in banana are for mature leaves. The two available samples suggest tentative concentration factors of 0.73 for ^{90}Sr , 0.42

for ^{137}Cs , and 0.0039 for $^{239,240}\text{Pu}$ in mature banana leaves. Unpublished data for banana fruit and soil samples from the same area collected in the 1972 Bikini survey yield concentration factors of < 0.058 for ^{90}Sr and < 0.00028 for $^{239,240}\text{Pu}$.¹⁷ Because no data are currently available for concentration factors for ^{137}Cs in bananas, we have assigned the conservative value of the ^{137}Cs concentration factor in mature banana leaves to banana fruit.

Messerschmidia and *Scaevola* — As discussed in the Enewetak survey,¹ the data from these two indicator plants show a significant correlation between leaves and soil on both an individual and a pooled basis. The number of samples from the Bikini survey does not warrant individual statistical analysis of these species. However, the pooled data show a correlation between leaves and soil that is significant at the 0.1 level for ^{90}Sr (Fig. 8) but is not significant for ^{137}Cs (Fig. 9). Cesium-137 does show a significant correlation between leaves and soil in the Enewetak results and probably would do so for Bikini if the number of samples were larger. Based on the median calculated concentration factors and on the results of the regression analysis, concentration factors of 0.49 for ^{90}Sr , 3.7 for ^{137}Cs , and 0.035 for

$^{239,240}\text{Pu}$ were assigned to mature *Scaevola* and *Messerschmidia* leaves.

Squash — One sample of summer squash and its seeds was available from the 1975 survey. This is the first time measured radionuclide concentrations in garden vegetables from Bikini Atoll have been available. Squash uptake of ^{90}Sr and ^{137}Cs is greater than that of any other plant sampled. Other workers also have observed high concentrations of ^{137}Cs in garden vegetables as compared to concentrations in other edible and indicator plants. Lynch *et al.*⁷ notes that in field studies, the ^{137}Cs concentration in lettuce leaves is an order of magnitude greater than the concentrations measured in other edible portions of food plants. In laboratory experiments to determine the uptake of ^{137}Cs by squash, Walker *et al.* report ^{137}Cs concentrations in squash that are higher than those measured in *Messerschmidia*, *Scaevola*, and *Pandanus* grown in the same Rongelap Atoll soil.¹⁸ However, from Walker's experiments it appears that with the application of fertilizer, the concentrations in squash can be reduced to levels comparable to those found in other edible plants.

Concentration Ratios

Radionuclides that are taken up from the soil through roots are either retained in the roots or transported

to the aboveground plant organs. At Bikini, the concentrations of ^{90}Sr in the aboveground plant organs of a species are comparable to the concentrations retained in the roots of that species. However, concentrations of ^{137}Cs and $^{239,240}\text{Pu}$ are lower in the aboveground plant organs than in roots (Table 11). In addition, there are differences in uptake among the various aboveground plant organs. Within any one species, concentrations of ^{90}Sr and $^{239,240}\text{Pu}$ are generally smaller in edible plant parts (e.g., fruit) than in nonedible organs (e.g.,

leaves); the opposite is true for ^{137}Cs (Table 12).

Because leaves are more often available for sampling than are fruit, we developed fruit-leaf concentration ratios to allow prediction of radio-nuclide concentrations in fruit from those measured in leaves of the same species (Table 13). The small number of samples makes it impossible to statistically evaluate these ratios; we will do this as more data become available. We also calculated leaf-leaf and fruit-fruit concentration ratios between different species for

Table 12. Summary of median soil-plant concentration factors.

| Species | Concentration Factor, (pCi/g dry plant)/(pCi/g dry soil) | | | | | |
|--|--|--------------------|-------------------|-------------------|-----------------------|---------------------|
| | ^{90}Sr | | ^{137}Cs | | $^{239,240}\text{Pu}$ | |
| | Mature Leaves | Fruit | Mature Leaves | Fruit | Mature Leaves | Fruit |
| Pooled <i>Scaevola</i> & <i>Messerschmidia</i> | 0.48 | --- | 3.7 | --- | 0.035 | --- |
| Coconut | 0.16 | 0.024 | 3.0 | 2.5 | 0.015 | --- |
| <i>Panadanus</i> | 0.91 | 0.50 | 15.0 | 5.4 | --- | --- |
| Papaya | 1.00 | 0.43 | 3.1 | 8.2 | 0.016 | 0.002 |
| Breadfruit | 1.80 | 0.76 | 1.6 | 7.0 | 0.027 | --- |
| Banana | 0.73 | 0.058 ^b | 0.42 | --- | 0.004 | 0.0003 ^b |
| Squash | 3.40 ^c | 0.15 ^d | 26.0 ^c | 56.0 ^d | --- | --- |

^a Not detected.

^b 1974 unpublished plant and soil data from the same vicinity.¹⁷

^c Whole plant.

^d Seeds.

Table 13. Fruit-leaf concentration ratios.

| Species | Nuclide | Concentration, pCi/g dry weight | | Fruit/Leaf Concentration Ratio |
|-----------------------|-------------------|---------------------------------|---------------|--------------------------------|
| | | Fruit | Mature Leaves | |
| <i>Pandanus</i> | ^{90}Sr | 40 ^a | 8 | 0.50 |
| | ^{137}Cs | 2.7 | 146 | 0.02 |
| Breadfruit | ^{90}Sr | 61 | 190 | 0.33 |
| | ^{137}Cs | 384 | 132 | 3.00 |
| Coconut "fruit" | ^{90}Sr | 0.79 | 12 | 0.06 ^c |
| | | 1.9 | 2.9 | |
| | ^{137}Cs | 60 | 7.1 | 4.00 ^c |
| | | 14 | 9.7 | |
| Coconut meat/milk | | 54 | 15 | |
| | ^{90}Sr | 0.41/0.18 | 1.7 | 0.33 |
| | ^{137}Cs | 76/9.3 | 24 | 3.00 |
| | | | | |
| Papaya | ^{90}Sr | 18 | 264 | 0.20 ^c |
| | | 22 ^a | 22 | |
| | | 49 | 191 | |
| | | 47 | 221 | |
| | ^{137}Cs | 865 ^b | 189 | 5.00 |
| | | 160 ^a | 156 | |
| | | 281 | 45 | |
| | | 303 | 69 | |
| $^{239,240}\text{Pu}$ | | 0.014 ^b | 0.049 | 0.03 |
| | | 0.0023 | 0.067 | |

^a Green fruit.^b Fallen fruit.^c Based on all available data.

prediction of concentrations in fruit of one species from those in fruit or leaves of another species. These ratios show that distribution patterns for each nuclide are consistent within a particular species. However, we must remember that the importance of the contribution of each nuclide to the internal dose to man varies with different species.

The fruit-fruit and fruit-leaf concentration ratios are calculated from a comparison of the concentration factors of three plant groups (plants within each group were sampled from the same general location) and from comparisons of the median concentration ratios of all associated plant-soil samples (Table 14). Analysis of the concentration ratios for mature leaves and fruit suggests that some species concentrate a given nuclide to a much greater extent than do others (Tables 15-19). For ^{90}Sr in mature leaves, the concentration decreases in the order: breadfruit and

papaya > *Pandanus* and banana > *Messerschmidia* and *Scaevola* > coconut (Tables 15 and 18). These results agree with those of Welander.¹⁰

The relative uptake of ^{137}Cs by the various species differs slightly from that of ^{90}Sr . For ^{137}Cs in mature leaves, the concentration appears to decrease with *Pandanus* > *Scaevola* > *Messerschmidia*, coconut, and papaya > breadfruit > banana. A comparison of ^{137}Cs uptake by fruit yields the pattern: papaya > *Pandanus* and breadfruit > coconut (Tables 16 and 19). The data for the uptake of $^{239,240}\text{Pu}$ by mature leaves are much more limited than data for ^{90}Sr and ^{137}Cs , but preliminary results suggest: *Messerschmidia* > breadfruit > *Pandanus* and coconut > papaya > *Scaevola* and banana (see Table 17). Although no concentration ratios are calculated, the uptake of ^{90}Sr and ^{137}Cs by unfertilized summer squash exceeds that of all other edible plants sampled.

Summary and Conclusions

The radionuclide concentration in surface soil samples (0 to 15 cm) varies greatly throughout both Bikini and Eneu Islands. In addition to the inhomogeneity observed in surface soil concentrations, profile data indicate that radionuclide concentration as a function of soil depth is quite variable. (In some cases,

the concentration at depths as great as 120 cm exceeds that in the top 2.5 cm.) As a result of the variability in surface soil concentrations with location and with depth, conclusions regarding dose reduction via soil removal must be exercised with great care. It is nearly impossible to generalize about remedial measures

Table 14. Associated soil-plant concentration factors for plant species sampled from the same location.

| Species, Organ | Concentration Factor, (pCi/g dry plant)/(pCi/g dry soil) | | |
|---------------------------------------|--|-------------------|------------------------|
| | ⁹⁰ Sr | ¹³⁷ Cs | ^{239, 240} Pu |
| <u>Group 1^a</u> | | | |
| Papaya, mature leaves | 1.6 | 3.9 | 0.012 |
| Papaya, mature leaves | 0.62 | 5.9 | 0.037 |
| Papaya, mature leaves | 0.94 | 0.30 | 0.0051 |
| Papaya, mature leaves | 4.0 | 2.9 | 0.027 |
| Banana, mature leaves | 0.48 | 0.33 | 0.0017 |
| Banana, mature leaves | 1.1 | 0.41 | 0.0060 |
| <u>Group 2^b</u> | | | |
| Breadfruit, mature leaves | 2.3 | 2.4 | 0.063 |
| <i>Pandanus</i> , mature leaves | 1.2 | 2.9 | 0.022 |
| <i>Messerschmidia</i> , mature leaves | 0.48 | 87 | 0.035 |
| <i>Messerschmidia</i> , mature leaves | 0.52 | 3.7 | --- |
| <i>Scaevola</i> , mature leaves | 0.24 | 1.3 | 0.0049 |
| <i>Scaevola</i> , mature leaves | 0.41 | 14 | --- |
| Breadfruit, fruit | 0.76 | 7.0 | --- |
| <i>Pandanus</i> , green fruit | 0.53 | 5.4 | --- |
| <i>Scaevola</i> , fruit | 0.14 | 1.4 | 0.00096 |
| <u>Group 3^c</u> | | | |
| <i>Scaevola</i> , mature leaves | 4.3 | 39 | 0.024 |
| Coconut, mature leaves | 0.67 | 14 | 0.026 |
| Coconut, mature leaves | 0.27 | 7.3 | 0.020 |

^a Group 1 includes samples T0010, T0030, T0040, T0050, T0060, T0070.

^b Group 2 includes samples T0090, T0100, T0110, T0120, and T0130.

^c Group 3 includes samples T0150 and T0160.

^d Not detected.

Table 15. Strontium-90 leaf-leaf concentration ratios.

| Species | Bread-fruit | Leaf-Leaf Concentration Ratio | | | | | |
|----------------|-------------|-------------------------------|----------|--------|-----------------|----------|---------|
| | | Papaya | Pandanus | Banana | Messer-schmidia | Scaevola | Coconut |
| Breadfruit | 1.0 | 1.5 | 2.0 | 2.5 | 3.5 | 6.0 | 12 |
| Papaya | 0.66 | 1.0 | 1.3 | 1.7 | 2.5 | 4.0 | 8.0 |
| Pandanus | 0.50 | 0.77 | 1.0 | 1.3 | 1.8 | 3.0 | 6.0 |
| Banana | 0.40 | 0.59 | 0.77 | 1.0 | 1.4 | 2.2 | 4.4 |
| Messerschmidia | 0.29 | 0.40 | 0.56 | 0.71 | 1.0 | 1.6 | 3.2 |
| Scaevola | 0.17 | 0.25 | 0.33 | 0.45 | 0.63 | 1.0 | 2.0 |
| Coconut | 0.08 | 0.12 | 0.17 | 0.23 | 0.31 | 0.50 | 1.0 |

Table 16. Cesium-137 leaf-leaf concentration ratios.

| Species | Leaf-Leaf Concentration Ratio | | | | | | Bread-fruit | Banana |
|----------------|-------------------------------|----------|-----------------|---------|--------|------|-------------|--------|
| | Pandanus | Scaevola | Messer-schmidia | Coconut | Papaya | | | |
| Pandanus | 1.0 | 2.0 | 4.0 | 5.0 | 5.0 | 10.0 | 35 | |
| Scaevola | 0.50 | 1.0 | 2.0 | 2.5 | 2.5 | 5.0 | 18 | |
| Messerschmidia | 0.25 | 0.50 | 1.0 | 1.3 | 1.3 | 2.4 | 8.8 | |
| Coconut | 0.20 | 0.40 | 0.80 | 1.0 | 1.0 | 2.0 | 7.0 | |
| Papaya | 0.20 | 0.40 | 0.80 | 1.0 | 1.0 | 2.0 | 7.5 | |
| Breadfruit | 0.10 | 0.20 | 0.40 | 0.50 | 0.50 | 1.0 | 3.8 | |
| Banana | 0.03 | 0.06 | 0.12 | 0.14 | 0.13 | 0.26 | 1.0 | |

entailing soil removal without first detailing the area and pathways that will be involved.

For example, the soil profile data (Appendices A and B) for Eneu and for some areas on Bikini indicate that removal of the top 10 cm of soil should do very little to reduce gamma exposure unless the removed soil is replaced with

clean soil, thus shielding the deeper contaminated soil. There would also be very little impact upon uptake by plants because the soil concentration is essentially identical through the root zone up to depths of 40 cm.

However, there are other areas on Bikini Island, (Appendices A and B, locations 501, 502, 503, and 504,

Table 17. Plutonium-239,240 leaf-leaf concentration ratios.

| Species | Leaf-Leaf Concentration Ratio | | | |
|------------------------------|-------------------------------|-------------------------|--------|------------------------------|
| | <i>Hesscherschmidia</i> | Breadfruit ^a | Papaya | <i>Seacvola</i> ^b |
| <i>Hesscherschmidia</i> | 1.0 | 4.5 | 10 | 15 |
| Breadfruit ^a | 0.22 | 1.0 | 2.2 | 3.3 |
| Papaya | 0.10 | 0.45 | 1.0 | 1.5 |
| <i>Seacvola</i> ^b | 0.07 | 0.30 | 0.67 | 1.0 |

^a Also includes *Pandanus* and coconut.^b Also includes banana.

Table 18. Strontium-90 fruit-fruit concentration ratios.

| Species | Fruit-Fruit Concentration Ratios | | |
|------------------------------|----------------------------------|------------------------------|---------|
| | Breadfruit | <i>Pandanus</i> ^a | Coconut |
| Breadfruit | 1.0 | 1.5 | 3.0 |
| <i>Pandanus</i> ^a | 0.67 | 1.0 | 2.0 |
| Coconut | 0.33 | 0.50 | 1.0 |

^a Also includes papaya.

Table 19. Cesium-137 fruit-fruit concentration ratios.

| Species | Fruit-Fruit Concentration Ratio | | | |
|-----------------|---------------------------------|------------|-----------------|---------|
| | Papaya | Breadfruit | <i>Pandanus</i> | Coconut |
| Papaya | 1.0 | 1.3 | 1.6 | 3.2 |
| Breadfruit | 0.75 | 1.0 | 1.4 | 2.8 |
| <i>Pandanus</i> | 0.63 | 0.71 | 1.0 | 2.0 |
| Coconut | 0.30 | 0.36 | 0.50 | 1.0 |

respectively) where removal of the top 10 cm of soil would reduce soil concentrations by approximately five-fold. This of course would result in reduced external exposure and reduced uptake in plants grown in such areas.

Although considerable variation in soil radionuclide concentration is observed on both islands, the soil concentrations on Bikini Island are approximately ten times those on Eneu Island. In addition, the relative radionuclide soil concentrations are very consistent; concentrations of ^{90}Sr and ^{137}Cs are ten to twenty times greater than those of $^{239,240}\text{Pu}$ and ^{241}Am which, in turn, are two to three times greater than ^{60}Co concentrations. Therefore, generalizations can safely be made from the soil data to the effect that inhabitants on Bikini Island will be exposed to higher doses than those on Eneu Island. Also, on both islands, ^{90}Sr and ^{137}Cs are the radionuclides of primary importance.

In the past, concentrations in terrestrial foodstuffs at Bikini Atoll have been predicted from soil concentrations measured in the field and from concentration factors taken from the literature. This approach was adopted because vegetation sampling programs were limited in the early surveys. However, the terrestrial sampling program of the 1975 Bikini survey included sufficient vegetation

samples to allow preliminary prediction of concentrations in nearly all components of the postulated Bikini diet. These predicted concentrations are based on soil-plant concentration factors and on fruit-leaf, leaf-leaf, or fruit-fruit concentration ratios calculated from the 1975 field data. The predicted concentrations compare favorably with the available measured concentrations. A more extensive survey of Bikini Atoll with larger sample sizes is needed to statistically verify these preliminary results. In the meantime, potential future concentrations in foodstuffs from Bikini can be predicted from our concentration factors if measured soil concentrations are available and from our concentration ratios if only vegetation samples are available. Estimates of the dose commitment expected from various projected lifestyles from our predicted concentrations are reported in Part 5 of this report series.⁹

The predominant nuclides in the terrestrial foodchain are ^{90}Sr and ^{137}Cs , followed by $^{239,240}\text{Pu}$, and will constitute the major internal dose to man from this pathway. In general, ^{137}Cs uptake by fruit and leaves is one order of magnitude greater than ^{90}Sr uptake which, in turn, exceeds $^{239,240}\text{Pu}$ uptake by one to two orders of magnitude. Uptake by mature leaves of papaya, banana, and breadfruit varies

5
0
1
0
1
3

slightly from this general pattern; in these cases, the measured uptake of ^{90}Sr is equal to that of ^{137}Cs . However, more samples are needed to verify the pervasiveness of these exceptions to the overall patterns observed. The distribution of ^{90}Sr , ^{137}Cs , and $^{239,240}\text{Pu}$ in fruit and leaves follows similar patterns in the various species studied. In a given species, ^{90}Sr and $^{239,240}\text{Pu}$ uptake by mature leaves is two to ten times greater than that by fruit. The ^{137}Cs uptake measured in this study shows a different trend;

uptake by fruit exceeds uptake by mature leaves by a factor of two to five.

A comparison of the uptake of ^{90}Sr , ^{137}Cs , and $^{239,240}\text{Pu}$ shows that, in general, the relative order of uptake is: squash (high; breadfruit, *Pandanus*, and papaya (intermediate); and banana and coconut (low). Slight variations occur, depending on the radionuclide under consideration. A more quantitative ordering can be made after greater numbers of all tissues of plants are sampled in future surveys.

Acknowledgment

We greatly appreciate the invaluable comments of Yook Ng and John Holdren in reviewing this manuscript. We thank John Koshiver for his aid in statistical analysis of the data and Ora Lowe and Bill Burke for their technical assistance.

The field portion of the June 1975 radiological survey of Bikini and Eneu Islands of Bikini Atoll was accomplished by a very intense and thorough effort of 21 people representing six different organizations. The number of samples collected and the amount of information obtained during the ten-day survey is a direct result of the cooperation and diligent effort of the following individuals: Wayne Bliss (Environmental Protection Agency, Las Vegas, Nevada), Bruce Clegg (Lawrence

Livermore Laboratory), Dave Coles (Lawrence Livermore Laboratory), Tom Crites (Lawrence Livermore Laboratory), Rod Eagle (Lawrence Livermore Laboratory), Harley Erwicker (Trust Territory of the Pacific Islands), Nat Greenhouse (Brookhaven National Laboratory), Paul Gudiksen (Lawrence Livermore Laboratory), Gale Holladay (Lawrence Livermore Laboratory), Bob Keller (Nevada Operations Office, ERDA), Dennis McBreen (Trust Territory of the Pacific Islands), Tommy McCraw (Division of Operational Safety, ERDA), Ben Mendoza (Lawrence Livermore Laboratory), Vic Nelson (University of Washington), Vic Noshkin (Lawrence Livermore Laboratory), Frank Reed (Environmental Protection Agency, Las Vegas, Nevada), Jim Schweiger (Lawrence Livermore Laboratory), Robert

Spies (Lawrence Livermore Laboratory), John Stewart (Nevada Operations Office, ERDA), and Marshall Stuart (Lawrence Livermore Laboratory).

We extend thanks to the following people for the superb job done in the processing of all soil and vegetation samples collected during the survey: Jim Schweiger (Lawrence Livermore Laboratory), Ben Mendoza (Lawrence Livermore Laboratory), Greg Calvaird (RECO), Elizabeth Fletcher (RECO), Nancy Sawley (Lawrence Livermore Laboratory), Cynthia Tafoya (Lawrence Livermore Laboratory), and Marshall Stuart (Lawrence Livermore Laboratory).

We are also grateful to the following people from McClellan Central Laboratory for the excellent effort in the wet chemistry analysis of the soil, vegetation, and animal samples: A. Ackland, R. Aduddel, L. Alexander, H. Aning, D. Beach, M. Beckinger, P. Carlson, W. Clark, R. Draper, W. Dunlap, D. Efurd, H. Erdman, D. Fletcher, W. Fuqua, C. Gay, J. Cholson, D. Griswold, R. Grogg, J. Hadl, H. Hamilton, R. Haslett, E. Henry, L. Hume, R. Jefferies, R. Johnson, M. Kantelo, P. Leciejewski, J. Lucas, R. Mayhew, C. McBrearty, G. Merrill, J. Miles, J. Miner, M. Mount, W. Myers, T. Nibarger, T. Opiela, R. Osborne, A. Paglione, J. Phelps, J. Phillips, C. Rheault,

J. Riggs, R. San Miguel, B. Scholl, R. Schwarting, D. Seymour, J. Sexton, P. Sparman, W. Summers, W. Tracey, R. Wagoner, W. Washer, L. Williams, and J. Wright.

We thank the following Lawrence Livermore Laboratory personnel involved in the gamma spectrometry of all the samples: Jesse Meadows, Mike Allen, Ruth Anderson, and Robert Wikkerink.

William Phillips has done a fine job in establishing, writing, updating, and correcting the rather extensive data bank generated by the analytical efforts.

The survey crew extends its thanks to Dr. Guy Haywood for medical support during the survey operation and to the Nevada Operations Office and Pacific Area Support Office for support services which enabled a smooth and efficient survey. Support from the Kwajalein Missile Range and the site contractor, Global Associates, as well as from the crew of the R.V. Liktanur is greatly appreciated.

The outstanding cooperation of personnel from the Trust Territory of the Pacific Islands and from the Office of the District Administrator of the Marshall Islands as well as that of the Bikini people played an important part in the successful completion of the survey.

References

1. *Enewetak Radiological Survey*, U.S. Atomic Energy Commission, Rept. NVO-140, vols. 1,2 (1973).
2. E. E. Held, *Radiological Resurvey of Animals, Soils and Groundwater at Bikini Atoll, 1969-70*, U.S. Atomic Energy Commission, Rept. NVO-269-8, Rev. 1 (1971).
3. P. A. Gustafson, *Radiological Report on Bikini Atoll*, Argonne National Laboratory Report (1968).
4. W. L. Robison, Lawrence Livermore Laboratory, private communication (1974).
5. P. H. Gudiksen, T. R. Crites, and W. L. Robison, *External Dose Estimates for Future Bikini Atoll Inhabitants*, Lawrence Livermore Laboratory, Rept. UCRL-51879, Part 1, Rev. 1 (1976).
6. V. E. Noshkin, W. L. Robison, K. M. Wong, and R. J. Eagle, *Preliminary Evaluation of the Radiological Quality of the Water on Bikini and Eneu Islands in 1975: Dose Assessment Based on Initial Sampling*, Lawrence Livermore Laboratory, Rept. UCRL-51879, Part 4 (1977).
7. O. D. T. Lynch, Jr., T. F. McCraw, W. A. Nelson, and W. E. Moore, *Radiochemical Resurvey of Food, Soil, Air and Groundwater at Bikini Atoll, 1972*, U. S. Energy Research and Development Administration, Publ. 34 (1975).
8. A. D. Welander, "Distribution of Radionuclides in the Environment of Enewetak and Bikini Atolls, August 1964," in D. J. Nelson and F. C. Guand, Eds., *Symposium on Radioecology*, U.S. Atomic Energy Commission, Rept. CONF-670503 (1969).
9. W. L. Robison, W. A. Phillips, and C. S. Colsher, *Dose Assessment at Bikini Atoll*, Lawrence Livermore Laboratory, Rept. UCRL-51879, Part 5 (1977).
10. J. F. Gamble and S. C. Snedaker, *Bioenvironmental and Radiological Feasibility Studies. Atlantic-Pacific Interceanic Canal Final Report - Agricultural Ecology*, Battelle Memorial Institute, Rept. BMI-821.02 (1969).
11. M. Mount, W. L. Robison, S. E. Thompson, K. O. Hamby, A. L. Prindle, and H. B. Levy, *Analytical Program - 1975 Bikini Radiological Survey*, Lawrence Livermore Laboratory, Rept. UCRL-51879, Part 2 (1976).
12. P. H. Gudiksen and O. D. T. Lynch, Jr., "Radioactivity Levels in Enewetak Soil," *Health Physics* 29, 17 (1975).
13. W. L. Robison and J. J. Koranda, Lawrence Livermore Laboratory, unpublished data (1976).
14. E. E. Held, S. P. Gessel, and R. B. Walter, *Atoll Soil Types in Relation to the Distribution of Fallout Radionuclides*, University of Washington, Laboratory of Radiation Biology, Rept. VWFL-92 (1965).

5010116 [REDACTED]

15. R. M. Kennedy, *The Soils of Rongelap Atoll*, M.S. thesis, University of Washington, Seattle, Washington (1962).
16. N. A. Greenhouse and T. F. McCraw, *Marshall Islands Radiological Followup*, Brookhaven National Laboratory, Rept. BNL-20767 (1976).
17. N. A. Greenhouse, Brookhaven National Laboratory, unpublished data (1974).
18. R. B. Walker, E. E. Held, and S. P. Gessel, "Radiocesium in Plants Grown on Rongelap Atoll Soils," *Recent Advances in Botany* 88, 1363 (1961).

PLL/gw/vt

BLANK PAGE

5010118

Appendix A. Soil Profiles of Bikini and Eneu Islands

The following tables present the concentration of eight selected radio-nuclides (^{60}Co , ^{90}Sr , ^{137}Cs , ^{155}Eu , ^{239}Pu , ^{240}Pu , ^{241}Pu , and ^{241}Am) with depth in the soil profile. Sample locations on the islands are given in Figs. 2 and 3. These data are also presented graphically in Appendix B.

PROFILE BIKINI 501

CONCENTRATIONS IN PCI/GM. MINUS FOR NO DATA. LESS THAN SIGN INDICATES INSTRUMENTAL DETECTION LIMIT

| DEPTH(CM) | 27 CO 60 | 38 SR 90 | 55 CS 137 | 63 EU 155 | 94 PU 239 | 94 PU 240 | 94 PU 241 | 95 AM 241 | MASTER LOG |
|-----------|-----------|-----------|------------|-----------|-----------|-----------|-----------|-------------|------------|
| 000-005 | 4.351E+00 | 2.370E+02 | 2.2260E+02 | 5.744E+00 | 1.311E+01 | 1.449E+01 | 2.607E+02 | 1.373E+01 | 01048131 |
| 005-010 | 5.869E+00 | 3.271E+02 | 1.365E+02 | 7.946E+00 | 1.698E+01 | 1.866E+01 | 3.387E+02 | 2.053E+01 | 01048234 |
| 010-020 | 3.853E+00 | 2.942E+02 | 8.473E+01 | 4.734E+00 | 1.112E+01 | 1.251E+01 | 2.330E+02 | 1.253E+01 | 01048371 |
| 020-030 | 6.261E-02 | 2.046E+02 | 3.219E+01 | 9.144E-02 | 6.748E-01 | 7.455E-01 | 1.312E+01 | 1.539E-01 | 01048472 |
| 030-040 | 9.230E-02 | 8.815E+01 | 1.305E+01 | 9.716E-02 | 4.586E-02 | 5.455E-02 | - | 1.9.032E-02 | 01048573 |

PROFILE BIKINI 502

CONCENTRATIONS IN PCI/GM. MINUS FOR NO DATA. LESS THAN SIGN INDICATES INSTRUMENTAL DETECTION LIMIT

| DEPTH(CM) | 27 CO 60 | 38 SR 90 | 55 CS 137 | 63 EU 155 | 94 PU 239 | 94 PU 240 | 94 PU 241 | 95 AM 241 | MASTER LOG |
|-----------|-----------|-----------|-----------|-------------|-----------|-----------|-----------|-------------|------------|
| 000-005 | 4.658E+00 | 2.871E+02 | 2.946E+02 | 7.270E+00 | - | - | - | - | 1.780E+01 |
| 005-010 | 5.477E+00 | 4.493E+02 | 3.130E+02 | 7.649E+00 | 2.193E+01 | 2.409E+01 | 4.346E+02 | 2.040E+01 | 01054634 |
| 010-020 | 8.122E-01 | 1.111E+02 | 4.146E+01 | 6.851E-01 | 1.770E+00 | 2.009E+00 | 3.883E+01 | 1.950E+00 | 01054771 |
| 020-030 | 9.194E-02 | 6.932E+00 | 2.365E+00 | 9.2.469E-02 | 3.724E-02 | 4.734E-02 | - | 1.4.260E-02 | 01054872 |

A-3
5010171

PROFILE BIKINI 503

CONCENTRATIONS IN PCI/GM. MINUS FOR NO DATA. LESS THAN SIGN INDICATES INSTRUMENTAL DETECTION LIMIT

| DEPTH(CM) | 27 CO 60 | 38 SR 90 | 55 CS 137 | 63 EU 155 | 94 PU 239 | 94 PU 240 | 94 PU 241 | 95 AM 241 | MASTER LOG |
|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-------------|------------|
| 000-005 | 1.045E+00 | 6.869E+01 | 8.617E+01 | 1.398E+00 | 3.327E+00 | 3.718E+00 | 7.527E+01 | 3.495E+00 | 01052331 |
| 005-010 | 8.748E-01 | 1.204E+02 | 7.378E+01 | 2.061E-01 | 9.770E-01 | 1.229E+00 | 2.556E+01 | 1.164E+00 | 01052434 |
| 010-020 | 1.426E-01 | 7.036E+01 | 9.719E+00 | 4.423E-02 | 1.212E-01 | 1.466E-01 | 2.700E+00 | 4.8.077E-02 | 01052571 |

PROFILE BIKINI 504

CONCENTRATIONS IN PCI/GM. MINUS FOR NO DATA. LESS THAN SIGN INDICATES INSTRUMENTAL DETECTION LIMIT

| DEPTH(CM) | 27 CO 60 | 38 SR 90 | 55 CS 137 | 63 EU 155 | 94 PU 239 | 94 PU 240 | 94 PU 241 | 95 AM 241 | MASTER LOG |
|-----------|-----------|-----------|-----------|------------|-----------|-----------|-----------|-------------|------------|
| 000-005 | 4.100E+00 | 2.208E+02 | 2.224E+02 | 5.545E+00 | 1.262E+01 | 1.396E+01 | 2.523E+02 | 1.446E+01 | 01049231 |
| 005-010 | 3.942E+00 | 2.458E+02 | 1.880E+02 | 5.514E+00 | 1.153E+01 | 1.297E+01 | 2.345E+02 | 1.435E+01 | 01049334 |
| 010-020 | 3.188E-01 | 2.826E+01 | 2.687E+01 | 1.696E-01 | 6.860E-01 | 7.982E-01 | 1.660E+01 | 6.306E-01 | 01049471 |
| 020-030 | 5.468E-02 | 7.973E+00 | 8.955E+00 | 4.2961E-02 | 1.236E-01 | 1.593E-01 | - | 4.5.721E-02 | 01049572 |
| 030-040 | 4.314E-02 | - | 5.532E+00 | 4.2473E-02 | - | - | - | 4.291E-02 | 01049673 |
| 040-050 | 4.314E-02 | - | 5.198E+00 | 4.3410E-02 | - | - | - | 4.5.550E-02 | 01049774 |

5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
249
250
251
252
253
254
255
256
257
258
259
259
260
261
262
263
264
265
266
267
268
269
269
270
271
272
273
274
275
276
277
278
279
279
280
281
282
283
284
285
286
287
288
289
289
290
291
292
293
294
295
296
297
298
299
299
300
301
302
303
304
305
306
307
308
309
309
310
311
312
313
314
315
316
317
318
319
319
320
321
322
323
324
325
326
327
328
329
329
330
331
332
333
334
335
336
337
338
339
339
340
341
342
343
344
345
346
347
348
349
349
350
351
352
353
354
355
356
357
358
359
359
360
361
362
363
364
365
366
367
368
369
369
370
371
372
373
374
375
376
377
378
379
379
380
381
382
383
384
385
386
387
388
389
389
390
391
392
393
394
395
396
397
398
399
399
400
401
402
403
404
405
406
407
408
409
409
410
411
412
413
414
415
416
417
418
419
419
420
421
422
423
424
425
426
427
428
429
429
430
431
432
433
434
435
436
437
438
439
439
440
441
442
443
444
445
446
447
448
449
449
450
451
452
453
454
455
456
457
458
459
459
460
461
462
463
464
465
466
467
468
469
469
470
471
472
473
474
475
476
477
478
479
479
480
481
482
483
484
485
486
487
488
489
489
490
491
492
493
494
495
496
497
498
499
499
500
501
502
503
504
505
506
507
508
509
509
510
511
512
513
514
515
516
517
518
519
519
520
521
522
523
524
525
526
527
528
529
529
530
531
532
533
534
535
536
537
538
539
539
540
541
542
543
544
545
546
547
548
549
549
550
551
552
553
554
555
556
557
558
559
559
560
561
562
563
564
565
566
567
568
569
569
570
571
572
573
574
575
576
577
578
579
579
580
581
582
583
584
585
586
587
588
589
589
590
591
592
593
594
595
596
597
598
599
599
600
601
602
603
604
605
606
607
608
609
609
610
611
612
613
614
615
616
617
618
619
619
620
621
622
623
624
625
626
627
628
629
629
630
631
632
633
634
635
636
637
638
639
639
640
641
642
643
644
645
646
647
648
649
649
650
651
652
653
654
655
656
657
658
659
659
660
661
662
663
664
665
666
667
668
669
669
670
671
672
673
674
675
676
677
678
679
679
680
681
682
683
684
685
686
687
688
689
689
690
691
692
693
694
695
696
697
698
699
699
700
701
702
703
704
705
706
707
708
709
709
710
711
712
713
714
715
716
717
718
719
719
720
721
722
723
724
725
726
727
728
729
729
730
731
732
733
734
735
736
737
738
739
739
740
741
742
743
744
745
746
747
748
749
749
750
751
752
753
754
755
756
757
758
759
759
760
761
762
763
764
765
766
767
768
769
769
770
771
772
773
774
775
776
777
778
779
779
780
781
782
783
784
785
786
787
788
789
789
790
791
792
793
794
795
796
797
798
799
799
800
801
802
803
804
805
806
807
808
809
809
810
811
812
813
814
815
816
817
818
819
819
820
821
822
823
824
825
826
827
828
829
829
830
831
832
833
834
835
836
837
838
839
839
840
841
842
843
844
845
846
847
848
849
849
850
851
852
853
854
855
856
857
858
859
859
860
861
862
863
864
865
866
867
868
869
869
870
871
872
873
874
875
876
877
878
879
879
880
881
882
883
884
885
886
887
888
889
889
890
891
892
893
894
895
896
897
898
899
899
900
901
902
903
904
905
906
907
908
909
909
910
911
912
913
914
915
916
917
918
919
919
920
921
922
923
924
925
926
927
928
929
929
930
931
932
933
934
935
936
937
938
939
939
940
941
942
943
944
945
946
947
948
949
949
950
951
952
953
954
955
956
957
958
959
959
960
961
962
963
964
965
966
967
968
969
969
970
971
972
973
974
975
976
977
978
979
979
980
981
982
983
984
985
986
987
988
989
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869

PROFILE BIKINI 507

CONCENTRATIONS IN PCI/GM. MINUS FOR NO DATA. LESS THAN SIGN INDICATES INSTRUMENTAL DETECTION LIMIT

| DEPTH(CM) | 27 CO 60 | 38 SR 90 | 55 CS 137 | 63 EU 155 | 94 PU 239 | 94 PU 240 | 94 PU 241 | 95 AM 241 | MASTER LOG |
|-----------|------------|-----------|-----------|------------|-----------|-----------|-----------|------------|------------|
| 000-005 | 3.411E+00 | 2.681E+02 | 1.337E+02 | 4.358E+00 | 1.000E+01 | 1.086E+01 | 1.960E+02 | 1.151E+01 | 01050331 |
| 005-010 | 1.583E+00 | 1.687E+02 | 8.410E+01 | 1.849E+00 | 4.368E+00 | 4.730E+00 | 8.851E+01 | 4.554E+00 | 01050434 |
| 010-020 | 1.180E-01 | 1.824E+01 | 2.149E+01 | ~5.153E-02 | 2.189E-01 | 2.571E-01 | - | 1.838E-01 | 01050571 |
| 020-030 | ~3.806E-02 | 8.901E-01 | 3.829E+00 | ~3.120E-02 | - | - | - | ~5.239E-02 | 01050672 |

PROFILE BIKINI 508

CONCENTRATIONS IN PCI/GM. MINUS FOR NO DATA. LESS THAN SIGN INDICATES INSTRUMENTAL DETECTION LIMIT

| DEPTH(CM) | 27 CO 60 | 38 SR 90 | 55 CS 137 | 63 EU 155 | 94 PU 239 | 94 PU 240 | 94 PU 241 | 95 AM 241 | MASTER LOG |
|-----------|-----------|-----------|-----------|------------|-----------|-----------|-----------|------------|------------|
| 000-005 | 2.768E+00 | 1.990E+02 | 1.113E+02 | 3.413E+00 | 7.554E+00 | 8.095E+00 | 1.468E+02 | 8.730E+00 | 01051431 |
| 005-010 | 3.582E+00 | 2.325E+02 | 1.344E+02 | 4.273E+00 | 9.514E+00 | 1.023E+01 | 1.886E+02 | 1.046E+01 | 01051534 |
| 010-020 | 3.898E+00 | 2.979E+02 | 1.395E+02 | 4.450E+00 | 9.802E+00 | 1.062E+01 | 1.900E+02 | 1.148E+01 | 01051671 |
| 020-030 | 5.626E+00 | 4.192E+02 | 1.489E+02 | 7.095E+00 | 1.492E+01 | 1.597E+01 | 2.886E+02 | 1.795E+01 | 01051772 |
| 030-040 | 2.150E+00 | 1.970E+02 | 7.590E+01 | 2.925E+00 | 7.036E+00 | 7.365E+00 | 1.359E+02 | 6.604E+00 | 01051873 |
| 040-050 | 1.673E-01 | 9.770E+01 | 1.122E+01 | ~4.595E-02 | 2.141E-01 | 2.634E-01 | 5.775E+00 | ~8.477E-02 | 01051974 |

51
50
49
48
47
46
45
44
43
42
41
40
39
38
37
36
35
34
33
32
31
30
29
28
27
26
25
24
23
22
21
20
19
18
17
16
15
14
13
12
11
10
9
8
7
6
5
4
3
2
1
0
PROFILE ENEU 801

CONCENTRATIONS IN PCI/GM. MINUS FOR NO DATA. LESS THAN SIGN INDICATES INSTRUMENTAL DETECTION LIMIT

| DEPTH(CM) | 27 CO 60 | 38 SR 90 | 55 CS 137 | 63 EU 155 | 94 PU 239 | 94 PU 240 | 94 PU 241 | 95 AM 241 | MASTER LOG |
|-----------|-----------|-----------|-----------|------------|-----------|-----------|-----------|------------|------------|
| 000-005 | 1.000E-01 | 4.200E+00 | 8.680E+00 | 1.623E-01 | 3.222E-01 | 3.506E-01 | 6.770E+00 | 2.783E-01 | 06088031 |
| 005-010 | 6.883E-02 | 3.518E+00 | 3.870E+00 | 7.050E-02 | 2.194E-01 | 2.392E-01 | - | 2.321E-01 | 06088134 |
| 010-020 | 1.415E-01 | 5.977E+00 | 4.374E+00 | 4.174E-01 | 8.347E-01 | 1.013E+00 | - | 7.667E-01 | 06088271 |
| 020-030 | 6.721E-02 | 2.405E+00 | 8.590E-01 | ~3.649E-02 | 1.694E-01 | 1.960E-01 | - | 1.122E-01 | 06088372 |
| 030-040 | 4.028E-02 | 2.245E+00 | 1.206E-01 | ~3.123E-02 | - | - | - | ~5.536E-02 | 06088473 |

9-V
PROFILE ENEU 802

CONCENTRATIONS IN PCI/GM. MINUS FOR NO DATA. LESS THAN SIGN INDICATES INSTRUMENTAL DETECTION LIMIT

| DEPTH(CM) | 27 CO 60 | 38 SR 90 | 55 CS 137 | 63 EU 155 | 94 PU 239 | 94 PU 240 | 94 PU 241 | 95 AM 241 | MASTER LOG |
|-----------|------------|-----------|-----------|------------|-----------|-----------|-----------|------------|------------|
| 000-005 | ~1.049E-02 | 1.376E+00 | 1.389E+00 | ~2.058E-02 | 2.554E-02 | 2.933E-02 | 7.396E-01 | ~3.750E-02 | 06090431 |
| 005-010 | ~2.291E-02 | 1.447E+00 | 1.280E+00 | ~2.386E-02 | 2.053E-02 | 1.977E-02 | - | ~3.769E-02 | 06090534 |
| 010-020 | ~8.991E-03 | 1.031E+00 | 1.236E+00 | ~2.459E-02 | 1.764E-02 | 1.949E-02 | 3.985E-01 | ~5.288E-02 | 06090671 |
| 020-030 | ~3.527E-02 | 6.838E-01 | 2.068E+00 | ~5.450E-02 | 6.005E-02 | 6.842E-02 | 1.354E+00 | ~9.734E-02 | 06090772 |
| 030-040 | 7.243E-02 | 9.779E-01 | 2.668E+00 | ~5.842E-02 | 1.198E-01 | 1.388E-01 | - | ~9.838E-02 | 06090873 |

PROFILE ENEU 803

CONCENTRATIONS IN PCI/GM. MINUS FOR NO DATA. LESS THAN SIGN INDICATES INSTRUMENTAL DETECTION LIMIT

| DEPTH(CM) | 27 CO 60 | 38 SR 90 | 55 CS 137 | 63 EU 155 | 94 PU 239 | 94 PU 240 | 94 PU 241 | 95 AM 241 | MASTER LOG |
|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|------------|
| 000-005 | 2.489E-01 | 1.692E+01 | 1.073E+01 | 4.559E-01 | 1.160E+00 | 1.250E+00 | 2.267E+01 | 1.021E+00 | 06094631 |
| 005-010 | 2.432E-01 | 1.571E+01 | 1.034E+01 | 4.613E-01 | 1.150E+00 | 1.237E+00 | 2.396E+01 | 9.847E-01 | 06094734 |
| 010-020 | 2.691E-01 | 2.397E+01 | 8.743E+00 | 6.631E-01 | 1.279E+00 | 1.433E+00 | - | 1.318E+00 | 06094871 |
| 020-030 | 3.052E-01 | 2.930E+01 | 9.946E+00 | 6.743E-01 | 1.559E+00 | 1.654E+00 | - | 1.290E+00 | 06094972 |
| 030-040 | 3.271E-01 | 3.192E+01 | 1.109E+01 | 7.995E-01 | 1.939E+00 | 2.099E+00 | 4.055E+01 | 2.090E+00 | 06095073 |
| 040-050 | 2.232E-01 | 1.747E+01 | 1.035E+01 | 5.473E-01 | 1.138E+00 | 1.249E+00 | 3.631E+01 | 1.255E+00 | 06095174 |

PROFILE HFH2

CONCENTRATIONS IN PCI/GM. MINUS FOR NO DATA. LESS THAN SIGN INDICATES INSTRUMENTAL DETECTION LIMIT

| DEPTH(CM) | 27 CO 60 | 38 SR 90 | 55 CS 137 | 63 EU 155 | 94 PU 239 | 94 PU 240 | 94 PU 241 | 95 AM 241 | MASTER LOG |
|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|------------|
| 000-005 | 1.116E-01 | 5.662E+00 | 2.991E+01 | 1.628E-01 | 3.957E-01 | 4.429E-01 | - | 4.267E-01 | 01056790 |
| 005-010 | 1.504E-01 | 6.041E+00 | 2.989E+01 | 1.928E-01 | 4.797E-01 | 5.338E-01 | 1.036E+01 | 5.622E-01 | 01056891 |
| 010-015 | 1.409E-01 | 5.333E+00 | 1.937E+01 | 1.912E-01 | 3.909E-01 | 4.464E-01 | 1.139E+01 | 4.100E-01 | 01056992 |
| 015-025 | 6.829E-02 | 6.243E+00 | 3.800E+00 | 1.177E-01 | 3.090E-01 | 3.637E-01 | 9.086E-01 | 2.738E-01 | 01057093 |
| 025-035 | 1.985E-01 | - | 3.576E+00 | 2.465E-01 | - | - | - | 6.279E-01 | 01057194 |
| 035-045 | 1.086E-01 | - | 2.969E+00 | 1.741E-01 | - | - | - | 2.814E-01 | 01057295 |

C
07/01/05
A-8
PROFILE HFH3

CONCENTRATIONS IN PCI/GM. MINUS FOR NO DATA. LESS THAN SIGN INDICATES INSTRUMENTAL DETECTION LIMIT

| DEPTH(CM) | 27 CO | 60 | 38 SR | 90 | 55 CS | 137 | 63 EU | 155 | 94 PU | 239 | 94 PU | 240 | 94 PU | 241 | 95 AM | 241 | MASTER LOG |
|-----------|-----------|----|-----------|----|-----------|-----|------------|-----|-----------|-----|-----------|-----|-----------|-----|------------|-----|------------|
| 000-005 | 1.895E-01 | | 1.735E+01 | | 1.162E+01 | | 2.850E-01 | | 6.937E-01 | | 7.545E-01 | | - | | 4.229E-01 | | 01031390 |
| 005-010 | 1.132E-01 | | 1.183E+01 | | 5.315E+00 | | 1.165E-01 | | 2.838E-01 | | 3.074E-01 | | 5.950E+00 | | 2.601E-01 | | 01031491 |
| 010-015 | 1.927E-01 | | 1.065E+01 | | 8.117E+00 | | 1.804E-01 | | 3.960E-01 | | 4.470E-01 | | - | | 3.490E-01 | | 01031592 |
| 015-025 | 1.309E+00 | | 7.730E+01 | | 6.653E+01 | | 1.892E+00 | | 4.743E+00 | | 5.293E+00 | | 1.113E+02 | | 4.577E+00 | | 01031693 |
| 035-045 | 1.070E-01 | | 1.352E+01 | | 8.400E-01 | | 2.966E-02 | | 6.319E-02 | | 7.293E-02 | | - | | 4.264E-02 | | 01031895 |
| 045-055 | 1.158E-01 | | - | | 3.307E-01 | | ~5.743E-02 | | - | | - | | - | | ~6.302E-02 | | 01031996 |

PROFILE HFH4

CONCENTRATIONS IN PCI/GM. MINUS FOR NO DATA. LESS THAN SIGN INDICATES INSTRUMENTAL DETECTION LIMIT

| DEPTH(CM) | 27 CO | 60 | 38 SR | 90 | 55 CS | 137 | 63 EU | 155 | 94 PU | 239 | 94 PU | 240 | 94 PU | 241 | 95 AM | 241 | MASTER LOG |
|-----------|------------|----|-----------|----|------------|-----|------------|-----|-----------|-----|-----------|-----|-----------|-----|------------|-----|------------|
| 000-005 | 8.838E-01 | | 6.500E+01 | | 4.275E+01 | | 8.599E-01 | | 2.024E+00 | | 2.272E+00 | | 5.622E+01 | | 2.331E+00 | | 01058390 |
| 005-010 | 4.177E-01 | | 3.671E+01 | | 1.656E+01 | | 2.847E-01 | | 9.162E-01 | | 1.086E+00 | | 2.071E+01 | | 1.079E+00 | | 01058491 |
| 010-015 | 1.708E-01 | | 1.649E+01 | | 5.072E+00 | | ~3.336E-02 | | 2.391E-01 | | 3.150E-01 | | 6.176E+00 | | 1.974E-01 | | 01058592 |
| 015-025 | 1.048E-01 | | 1.809E+01 | | 2.542E+00 | | ~2.357E-02 | | 1.740E-01 | | 2.317E-01 | | - | | 1.860E-01 | | 01058693 |
| 025-035 | ~2.822E-02 | | - | | ~2.429E-02 | | ~1.896E-02 | | - | | - | | - | | ~3.231E-02 | | 01058794 |
| 035-045 | ~1.782E-02 | | - | | 2.241E-01 | | ~3.842E-02 | | - | | - | | - | | ~5.667E-02 | | 01058895 |

PROFILE HFH5

CONCENTRATIONS IN PCI/GM. MINUS FOR NO DATA. LESS THAN SIGN INDICATES INSTRUMENTAL DETECTION LIMIT

| DEPTH(CM) | 27 CO | 60 | 38 SR | 90 | 55 CS | 137 | 63 EU | 155 | 94 PU | 239 | 94 PU | 240 | 94 PU | 241 | 95 AM | 241 | MASTER LOG |
|-----------|------------|----|-----------|----|-----------|-----|------------|-----|-----------|-----|-----------|-----|-----------|-----|------------|-----|------------|
| 000-005 | 1.854E+00 | | 1.149E+02 | | 8.162E+01 | | 2.752E+00 | | 6.545E+00 | | 7.162E+00 | | 1.287E+02 | | 5.707E+00 | | 01029890 |
| 005-010 | 7.423E+00 | | 3.219E+02 | | 1.401E+02 | | 1.283E+01 | | 3.202E+01 | | 3.351E+01 | | 5.959E+02 | | 2.841E+01 | | 01029991 |
| 010-015 | 3.173E+00 | | 2.350E+02 | | 1.284E+02 | | 4.301E+00 | | 1.031E+01 | | 1.117E+01 | | 2.867E+02 | | 9.477E+00 | | 01030092 |
| 015-025 | 6.541E-02 | | 4.793E+01 | | 4.761E+00 | | ~2.122E-02 | | 4.883E-02 | | 5.730E-02 | | 1.162E+00 | | ~3.274E-02 | | 01030193 |
| 025-035 | ~3.274E-02 | | - | | 4.241E-01 | | ~2.227E-02 | | - | | - | | - | | ~3.768E-02 | | 01030294 |
| 035-045 | ~3.041E-02 | | - | | 4.824E-01 | | ~2.189E-02 | | - | | - | | - | | ~3.730E-02 | | 01030496 |

PROFILE TS0001

CONCENTRATIONS IN PCI/GM. MINUS FOR NO DATA. LESS THAN SIGN INDICATES INSTRUMENTAL DETECTION LIMIT

| DEPTH(CM) | 27 CO | 60 | 38 SR | 90 | 55 CS | 137 | 63 EU | 155 | 94 PU | 239 | 94 PU | 240 | 94 PU | 241 | 95 AM | 241 | MASTER LOG |
|-----------|------------|----|-----------|----|-----------|-----|------------|-----|-----------|-----|-----------|-----|-----------|-----|------------|-----|------------|
| 000-005 | 8.829E-01 | | 6.338E+01 | | 5.414E+01 | | 1.562E+00 | | 3.391E+00 | | 3.683E+00 | | 6.572E+01 | | 4.173E+00 | | 01000190 |
| 005-010 | 8.554E-01 | | 6.230E+01 | | 4.883E+01 | | 1.381E+00 | | 3.114E+00 | | 3.478E+00 | | 8.288E+01 | | 3.548E+00 | | 01000291 |
| 010-015 | 1.176E+00 | | 8.554E+01 | | 4.676E+01 | | 1.814E+00 | | 4.206E+00 | | 4.667E+00 | | 8.613E+01 | | 5.239E+00 | | 01000392 |
| 015-020 | 1.814E+00 | | 1.175E+02 | | 4.770E+01 | | 2.951E+00 | | 6.464E+00 | | 7.198E+00 | | 1.327E+02 | | 6.604E+00 | | 01000401 |
| 020-025 | 1.463E+00 | | 8.770E+01 | | 3.266E+01 | | 1.851E+00 | | 4.923E+00 | | 5.131E+00 | | 9.563E+01 | | 4.069E+00 | | 01000501 |
| 025-030 | 2.264E+00 | | 1.201E+02 | | 2.271E+01 | | 3.908E+00 | | 8.928E+00 | | 9.550E+00 | | 1.762E+02 | | 7.928E+00 | | 01000601 |
| 030-040 | ~3.828E-02 | | 3.669E+01 | | 8.041E+00 | | ~6.559E-02 | | 6.198E-02 | | 7.230E-02 | | - | | ~1.328E-01 | | 01000701 |
| 040-045 | - | | 2.453E+01 | | - | | - | | 6.743E-02 | | 9.185E-02 | | - | | - | | 01000801 |

6-V

50100
171022

PROFILE TS0002

CONCENTRATIONS IN PC1/GM. MINUS FOR NO DATA. LESS THAN SIGN INDICATES INSTRUMENTAL DETECTION LIMIT

| DEPTH(CM) | 27 CO 60 | 38 SR 90 | 55 CS 137 | 63 EU 155 | 94 PU 239 | 94 PU 240 | 94 PU 241 | 95 AM 241 | MASTER LOG |
|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|------------|
| 000-005 | 9.536E-01 | 4.797E+01 | 4.923E+01 | 1.385E+00 | 3.344E+00 | 3.672E+00 | 6.820E+01 | 2.979E+00 | 01006301 |
| 005-010 | 9.221E-01 | 5.743E+01 | 3.216E+01 | 1.187E+00 | 3.257E+00 | 3.642E+00 | 6.982E+01 | 2.749E+00 | 01006401 |
| 010-015 | 8.777E-01 | 6.022E+01 | 1.780E+01 | 1.383E+00 | 3.398E+00 | 3.778E+00 | 7.003E+01 | 3.202E+00 | 01006501 |
| 015-025 | 2.162E+00 | 2.015E+02 | 2.724E+01 | 3.900E+00 | 7.626E+00 | 8.378E+00 | 1.528E+02 | 8.122E+00 | 01006601 |
| 025-035 | 2.447E-01 | 6.541E+01 | 3.598E+00 | 2.014E-01 | 4.905E-01 | 5.252E-01 | 1.194E+01 | 4.284E-01 | 01006701 |

PROFILE TS0003

CONCENTRATIONS IN PC1/GM. MINUS FOR NO DATA. LESS THAN SIGN INDICATES INSTRUMENTAL DETECTION LIMIT

| DEPTH(CM) | 27 CO 60 | 38 SR 90 | 55 CS 137 | 63 EU 155 | 94 PU 239 | 94 PU 240 | 94 PU 241 | 95 AM 241 | MASTER LOG |
|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|------------|
| 000-005 | 5.748E-01 | 4.389E+01 | 5.568E+01 | 8.554E-01 | 2.042E+00 | 2.300E+00 | 3.881E+01 | 1.818E+00 | 01106701 |
| 005-010 | 5.914E-01 | 4.040E+01 | 4.644E+01 | 8.811E-01 | 2.319E+00 | 2.543E+00 | 4.743E+01 | 1.814E+00 | 01106801 |
| 010-015 | 1.779E+00 | 9.928E+01 | 5.802E+01 | 3.255E+00 | 6.090E+00 | 6.779E+00 | 1.222E+02 | 6.946E+00 | 01006901 |
| 015-025 | 3.173E+00 | 2.433E+02 | 3.406E+01 | 4.352E+00 | 1.024E+01 | 1.127E+01 | 2.538E+02 | 9.432E+00 | 01007001 |
| 025-035 | 1.142E+00 | 1.400E+02 | 3.318E+01 | 1.321E+00 | 3.831E+00 | 3.981E+00 | 7.662E+01 | 2.945E+00 | 01007101 |
| 035-045 | - | 3.404E+01 | - | - | 4.277E-01 | 5.050E-01 | - | - | 01007201 |

PROFILE TS0012

CONCENTRATIONS IN PCI/GM. MINUS FOR NO DATA. LESS THAN SIGN INDICATES INSTRUMENTAL DETECTION LIMIT

| DEPTH(CM) | 27 CO 60 | 38 SR 90 | 55 CS 137 | 63 EU 155 | 94 PU 239 | 94 PU 240 | 94 PU 241 | 95 AM 241 | MASTER LOG |
|-----------|-----------|-----------|-----------|------------|-----------|-----------|-----------|------------|------------|
| 000-005 | 1.830E+00 | 1.911E+02 | 1.243E+02 | 2.649E+00 | 6.595E+00 | 7.293E+00 | 1.397E+02 | 5.464E+00 | 01010901 |
| 005-010 | 2.324E+00 | 1.961E+02 | 1.021E+02 | 3.145E+00 | 7.410E+00 | 8.203E+00 | 1.574E+02 | 8.811E+00 | 01011101 |
| 015-020 | 2.381E+00 | 2.034E+02 | 9.572E+01 | 3.791E+00 | 9.176E+00 | 9.964E+00 | 1.851E+02 | 1.060E+01 | 01011201 |
| 020-033 | 9.140E-01 | 8.869E+01 | 4.946E+00 | 1.301E+00 | 2.948E+00 | 3.395E+00 | 7.964E+01 | 2.772E+00 | 01011301 |
| 036-051 | 1.062E+00 | 3.014E+02 | 9.955E+01 | 9.631E-01 | 2.460E+00 | 2.671E+00 | 4.856E+01 | 2.131E+00 | 01011401 |
| 056-102 | 1.800E-02 | 1.207E+00 | 1.742E+00 | ±1.035E-02 | 1.338E-02 | 1.500E-02 | - | ±1.667E-02 | 01011501 |

PROFILE TS0031

CONCENTRATIONS IN PCI/GM. MINUS FOR NO DATA. LESS THAN SIGN INDICATES INSTRUMENTAL DETECTION LIMIT

| DEPTH(CM) | 27 CO 60 | 38 SR 90 | 55 CS 137 | 63 EU 155 | 94 PU 239 | 94 PU 240 | 94 PU 241 | 95 AM 241 | MASTER LOG |
|-----------|------------|------------|-----------|------------|-----------|-----------|-----------|------------|------------|
| 000-005 | 2.331E-01 | 2.757E+01 | 2.193E+01 | 2.991E-01 | 5.833E-01 | 6.883E-01 | 1.429E+01 | 7.045E-01 | 01004590 |
| 005-010 | 6.568E-01 | 5.527E+01 | 4.224E+01 | 5.595E-01 | 1.606E+00 | 1.845E+00 | 3.701E+01 | 1.945E+00 | 01004691 |
| 010-015 | 2.483E-01 | 3.643E+01 | 2.295E+01 | 1.774E-01 | 5.149E-01 | 5.712E-01 | - | 5.297E-01 | 01004792 |
| 015-025 | 1.741E-02 | 5.081E-01 | 1.498E+00 | ±1.814E-02 | - | - | - | ±2.913E-02 | 01004893 |
| 025-035 | ±2.910E-02 | - | 1.262E-01 | ±1.509E-02 | - | - | - | ±2.558E-02 | 01004994 |
| 035-045 | ±2.298E-02 | ±9.009E-01 | 1.103E-01 | ±1.581E-02 | 8.491E-04 | 7.667E-04 | - | ±2.886E-02 | 01005095 |

PROFILE TS0041

CONCENTRATIONS IN PCI/GM. MINUS FOR NO DATA. LESS THAN SIGN INDICATES INSTRUMENTAL DETECTION LIMIT

| DEPTH(CM) | 27 CO 60 | 38 SR 90 | 55 CS 137 | 63 EU 155 | 94 PU 239 | 94 PU 240 | 94 PU 241 | 95 AM 241 | MASTER LOG |
|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|------------|
| 000-005 | 2.516E-01 | 1.973E+01 | 1.734E+01 | 2.541E-01 | 6.761E-01 | 7.626E-01 | - | 6.126E-01 | 01002090 |
| 005-010 | 1.127E+00 | 4.950E+01 | 6.401E+01 | 1.318E+00 | 3.278E+00 | 3.691E+00 | 6.671E+01 | 2.936E+00 | 01002191 |
| 010-015 | 1.270E+00 | 3.368E+01 | 7.009E+01 | 1.242E+00 | 3.627E+00 | 4.234E+00 | - | 2.983E+00 | 01002292 |
| 015-025 | 2.064E-01 | 4.088E+01 | 1.295E+01 | 7.730E-02 | 9.041E-02 | 1.130E-01 | - | 7.106E-01 | 01002393 |
| 025-035 | 7.041E-02 | 5.653E+00 | 1.022E+01 | 7.092E-02 | 3.256E-02 | 3.532E-02 | - | 7.505E-02 | 01002494 |
| 035-045 | 7.441E-02 | 7.909E-01 | 4.104E+00 | 2.522E-02 | 4.905E-03 | 6.302E-03 | - | 7.990E-02 | 01002595 |

A-12

PROFILE TS0051

CONCENTRATIONS IN PCI/GM. MINUS FOR NO DATA. LESS THAN SIGN INDICATES INSTRUMENTAL DETECTION LIMIT

| DEPTH(CM) | 27 CO 60 | 38 SP 90 | 55 CS 137 | 63 EU 155 | 94 PU 239 | 94 PU 240 | 94 PU 241 | 95 AM 241 | MASTER LOG |
|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|------------|
| 000-005 | 2.416E+00 | 1.950E+02 | 1.229E+02 | 3.636E+00 | 7.878E+00 | 8.770E+00 | - | 6.892E+00 | 01003590 |
| 005-010 | 2.932E+00 | 2.032E+02 | 1.410E+02 | 3.870E+00 | 9.477E+00 | 1.050E+01 | 2.013E+02 | 8.329E+00 | 01003691 |
| 010-015 | 3.166E+00 | 2.420E+02 | 1.437E+02 | 4.426E+00 | 9.973E+00 | 1.117E+01 | - | 8.905E+00 | 01003792 |
| 015-025 | 3.004E+00 | 1.882E+02 | 1.748E+02 | 4.216E+00 | 9.302E+00 | 1.043E+01 | - | 9.054E+00 | 01003893 |
| 025-035 | 3.200E+00 | 2.097E+02 | 1.907E+02 | 4.432E+00 | 9.622E+00 | 1.068E+01 | - | 8.797E+00 | 01003994 |
| 035-045 | 2.865E+00 | 2.186E+02 | 1.824E+02 | 3.988E+00 | 9.288E+00 | 1.036E+01 | 2.137E+02 | 8.324E+00 | 01004095 |

PROFILE TS0061

CONCENTRATIONS IN PCI/GM. MINUS FOR NO DATA. LESS THAN SIGN INDICATES INSTRUMENTAL DETECTION LIMIT

| DEPTH(CM) | 27 CO 50 | 38 SR 90 | 55 CS 137 | 63 EU 155 | 94 PU 239 | 94 PU 240 | 94 PU 241 | 95 AM 241 | MASTER LOG |
|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|------------|
| 000-005 | 2.722E-01 | 5.122E+01 | 3.674E+01 | 5.059E-01 | 1.031E+00 | 1.142E+00 | 2.340E+01 | 9.284E-01 | 01008790 |
| 005-010 | 1.588E-01 | 1.769E+01 | 1.171E+01 | 3.266E-01 | 6.511E-01 | 7.159E-01 | 1.760E+01 | 7.197E-01 | 01008891 |
| 010-015 | 7.739E-02 | 7.518E+00 | 6.360E+00 | 1.100E-01 | 2.480E-01 | 2.801E-01 | - | 2.527E-01 | 01008992 |
| 015-025 | 1.855E+00 | 1.350E+02 | 8.252E+01 | 2.759E+00 | 6.198E+00 | 6.829E+00 | 1.294E+02 | 6.014E+00 | 01009093 |
| 025-035 | 4.626E+00 | 4.087E+02 | 2.190E+02 | 6.743E+00 | 1.657E+01 | 1.831E+01 | 3.330E+02 | 1.336E+01 | 01009194 |
| 035-045 | 5.604E+00 | 4.761E+02 | 3.393E+02 | 8.369E+00 | 1.729E+01 | 1.909E+01 | 3.495E+02 | 1.794E+01 | 01009295 |
| 045-055 | 3.285E+00 | 4.311E+02 | 1.350E+02 | 5.432E+00 | 1.104E+01 | 1.217E+01 | 2.263E+02 | 1.159E+01 | 01009396 |
| 055-065 | 3.212E-01 | 2.700E+02 | 2.432E+01 | 7.117E-02 | 2.686E-01 | 2.943E-01 | - | 1.161E-01 | 01009497 |
| 065-075 | 8.162E-02 | 5.324E+01 | 1.571E+00 | 2.708E-02 | 8.563E-03 | 1.001E-02 | 1.787E-01 | 1.875E-02 | 01009598 |

A-13

PROFILE TS0062

CONCENTRATIONS IN PCI/GM. MINUS FOR NO DATA. LESS THAN SIGN INDICATES INSTRUMENTAL DETECTION LIMIT

| DEPTH(CM) | 27 CO 50 | 38 SR 90 | 55 CS 137 | 63 EU 155 | 94 PU 239 | 94 PU 240 | 94 PU 241 | 95 AM 241 | MASTER LOG |
|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|------------|
| 000-005 | 8.162E-01 | 7.838E+01 | 5.833E+01 | 1.091E+00 | 2.539E+00 | 2.812E+00 | 5.252E+01 | 3.146E+00 | 01001190 |
| 005-010 | 5.387E-01 | 3.867E+01 | 2.744E+01 | 7.554E-01 | 1.744E+00 | 2.008E+00 | - | 1.979E+00 | 01001291 |
| 010-015 | 1.159E-01 | 7.892E+01 | 3.134E+00 | 4.191E-01 | 6.077E-01 | 6.563E-01 | 1.196E+01 | 5.081E-01 | 01001392 |
| 015-025 | 1.329E+00 | 7.622E+01 | 7.694E+01 | 2.023E+00 | 4.689E+00 | 5.167E+00 | 9.694E+01 | 4.054E+00 | 01001493 |
| 025-035 | 1.577E+00 | 1.493E+02 | 1.354E+02 | 1.767E+00 | 4.703E+00 | 5.347E+00 | - | 4.779E+00 | 01001594 |
| 035-045 | 1.575E-01 | 6.608E+01 | 1.616E+01 | 8.284E-02 | 7.842E-02 | 1.021E-01 | - | 1.245E-01 | 01001695 |

PROFILE TS0071

CONCENTRATIONS IN PCI/GM. MINUS FOR NO DATA. LESS THAN SIGN INDICATES INSTRUMENTAL DETECTION LIMIT

| DEPTH(CM) | 27 CO | 60 | 38 SR | 90 | 55 CS | 137 | 63 EU | 155 | 94 PU | 239 | 94 PU | 240 | 94 PU | 241 | 95 AM | 241 | MASTER LOG |
|-----------|-----------|-----------|-------|----|-----------|-----------|-----------|-----------|-----------|-----|-----------|-----|----------|-----|-------|-----|------------|
| 000-005 | 1.074E+00 | 1.071E+02 | | | 6.622E+01 | 1.412E+00 | 3.150E+00 | 3.594E+00 | 7.365E+01 | | 3.610E+00 | | 01005590 | | | | |
| 005-010 | 1.238E+00 | 1.140E+02 | | | 6.734E+01 | 1.600E+00 | 3.971E+00 | 4.382E+00 | 7.892E+01 | | 4.716E+00 | | 01005691 | | | | |
| 010-015 | 1.123E+00 | 1.046E+02 | | | 6.018E+01 | 1.667E+00 | 3.486E+00 | 3.886E+00 | 7.225E+01 | | 4.079E+00 | | 01006292 | | | | |
| 015-025 | - | 1.686E+02 | | | - | - | 6.662E+00 | 7.455E+00 | 1.429E+02 | | 7.779E+00 | | 01005792 | | | | |
| 025-035 | - | 7.943E+01 | | | - | - | 2.800E-01 | 3.616E-01 | 8.450E+00 | | - | | 01005893 | | | | |
| 035-045 | - | 4.963E+00 | | | - | - | 3.534E-02 | 4.308E-02 | 1.492E+00 | | - | | 01005994 | | | | |

PROFILE TS0081

CONCENTRATIONS IN PCI/GM. MINUS FOR NO DATA. LESS THAN SIGN INDICATES INSTRUMENTAL DETECTION LIMIT

| DEPTH(CM) | 27 CO | 60 | 38 SR | 90 | 55 CS | 137 | 63 EU | 155 | 94 PU | 239 | 94 PU | 240 | 94 PU | 241 | 95 AM | 241 | MASTER LOG |
|-----------|-----------|-----------|-------|----|-----------|------------|-----------|-----------|-----------|-----|------------|-----|----------|-----|-------|-----|------------|
| 000-005 | 6.707E-01 | 5.977E+01 | | | 3.821E+01 | 1.008E+00 | 1.952E+00 | 2.207E+00 | 4.640E+01 | | 2.355E+00 | | 01007490 | | | | |
| 005-010 | 2.109E+00 | 1.461E+02 | | | 8.149E+01 | 3.045E+00 | 7.108E+00 | 8.050E+00 | 1.769E+02 | | 8.775E+00 | | 01007591 | | | | |
| 010-015 | 2.968E+00 | 3.504E+02 | | | 1.036E+02 | 3.758E+00 | 1.036E+01 | 1.161E+01 | 2.409E+02 | | 1.334E+01 | | 01007692 | | | | |
| 015-025 | 3.510E-01 | 1.012E+02 | | | 2.183E+01 | 2.151E-01 | 5.883E-01 | 7.090E-01 | 1.410E+01 | | 6.072E-01 | | 01007793 | | | | |
| 025-035 | 7.757E-02 | 4.752E+01 | | | 6.144E+00 | ~3.423E-02 | 9.982E-02 | 1.282E-01 | - | | ~6.086E-02 | | 01007894 | | | | |
| 035-045 | 9.833E-02 | 2.414E+01 | | | 2.506E+00 | ~1.721E-02 | 7.910E-02 | 1.058E-01 | - | | 8.779E-02 | | 01007995 | | | | |
| 105-115 | 1.275E+00 | 9.991E+01 | | | 5.613E+01 | 2.305E+00 | 4.073E+00 | 4.391E+00 | 8.239E+01 | | 4.354E+00 | | 01008601 | | | | |

A-14

5010132

PROFILE TS0091

CONCENTRATIONS IN PCI/GM. MINUS FOR NO DATA. LESS THAN SIGN INDICATES INSTRUMENTAL DETECTION LIMIT

| DEPTH(CM) | 27 CO 60 | 38 SR 90 | 55 CS 137 | 63 EU 155 | 94 PU 239 | 94 PU 240 | 94 PU 241 | 95 AM 241 | MASTER LOG |
|-----------|------------|-----------|-----------|------------|-----------|-----------|-----------|------------|------------|
| 000-005 | 7.054E-01 | 4.389E+01 | 7.604E+01 | 1.126E+00 | 2.590E+00 | 2.943E+00 | 7.986E+01 | 3.207E+00 | 01028690 |
| 005-010 | 1.547E+00 | 8.883E+01 | 9.464E+01 | 1.925E+00 | 5.333E+00 | 5.964E+00 | 1.075E+02 | 6.054E+00 | 01028791 |
| 010-015 | 2.241E+00 | 1.301E+02 | 8.869E+01 | 3.482E+00 | 8.207E+00 | 9.212E+00 | 2.026E+02 | 9.464E+00 | 01028892 |
| 015-025 | 4.847E-01 | 8.356E+01 | 2.818E+01 | 3.001E-01 | 9.059E-01 | 1.073E+00 | - | 9.955E-01 | 01028993 |
| 025-035 | 1.619E-01 | 5.275E+01 | 7.072E+00 | ~7.311E-02 | 2.614E-01 | 3.117E-01 | 6.874E+00 | 2.877E-01 | 01029094 |
| 035-045 | ~3.901E-02 | 6.243E+00 | 2.636E+00 | ~2.936E-02 | 2.947E-02 | 3.685E-02 | - | ~4.653E-02 | 01029195 |
| 045-055 | ~2.391E-02 | - | 1.309E+00 | ~2.023E-02 | - | - | - | ~3.097E-02 | 01029296 |

PROFILE TS0101

CONCENTRATIONS IN PCI/GM. MINUS FOR NO DATA. LESS THAN SIGN INDICATES INSTRUMENTAL DETECTION LIMIT

| DEPTH(CM) | 27 CO 60 | 38 SR 90 | 55 CS 137 | 63 EU 155 | 94 PU 239 | 94 PU 240 | 94 PU 241 | 95 AM 241 | MASTER LOG |
|-----------|------------|-----------|------------|------------|-----------|-----------|-----------|------------|------------|
| 000-005 | 3.694E-01 | 7.905E+01 | 5.640E+01 | 4.937E-01 | 1.185E+00 | 1.319E+00 | 2.447E+01 | 1.374E+00 | 01035290 |
| 005-010 | 2.548E-01 | 4.991E+01 | 3.317E+01 | 2.854E-01 | 7.279E-01 | 8.441E-01 | 2.003E+01 | 7.991E-01 | 01035391 |
| 010-015 | 1.422E-01 | 3.244E+01 | 1.438E+01 | 7.946E-02 | 2.578E-01 | 2.864E-01 | 4.959E+00 | 2.747E-01 | 01035492 |
| 015-025 | ~3.252E-02 | 1.850E+01 | ~1.404E-01 | ~3.933E-02 | - | - | - | ~6.144E-02 | 01035593 |
| 025-035 | ~3.123E-02 | 6.302E+00 | 8.468E-01 | ~2.423E-02 | 1.060E-02 | 1.334E-02 | - | ~4.078E-02 | 01035694 |
| 035-045 | ~2.829E-02 | 3.993E+00 | 2.062E-01 | ~2.292E-02 | - | - | - | ~3.812E-02 | 01035795 |

ST-V

C110133

!

PROFILE TS0111

CONCENTRATIONS IN PCI/GM. MINUS FOR NO DATA. LESS THAN SIGN INDICATES INSTRUMENTAL DETECTION LIMIT

| DEPTH(CM) | 27 CO 60 | 38 SR 90 | 55 CS 137 | 63 EU 155 | 94 PU 239 | 94 PU 240 | 94 PU 241 | 95 AM 241 | MASTER LOG |
|-----------|------------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|------------|
| 000-005 | 6.153E-01 | 6.396E+01 | 7.946E+01 | 7.667E-01 | 2.147E+00 | 2.385E+00 | 5.162E+01 | 2.677E+00 | 01011990 |
| 005-010 | 9.784E-01 | 9.523E+01 | 9.968E+01 | 1.637E+00 | 3.474E+00 | 3.869E+00 | 7.977E+01 | 4.039E+00 | 01012091 |
| 010-015 | 1.600E+00 | 8.194E+01 | 1.271E+02 | 2.015E+00 | 5.356E+00 | 5.973E+00 | 1.143E+02 | 6.631E+00 | 01012192 |
| 015-025 | 1.074E+00 | 8.338E+01 | 7.086E+01 | 1.038E+00 | 3.150E+00 | 3.577E+00 | - | 2.554E+00 | 01012293 |
| 025-035 | 3.204E-02 | 3.616E+01 | 3.063E+00 | 1.523E-02 | 2.776E-02 | 3.598E-02 | - | 1.155E-02 | 01012394 |
| 035-045 | 1.2958E-02 | 7.306E+00 | 2.603E+00 | 1.286E-02 | - | - | - | 1.926E-02 | 01012495 |

PROFILE TS0121

CONCENTRATIONS IN PCI/GM. MINUS FOR NO DATA. LESS THAN SIGN INDICATES INSTRUMENTAL DETECTION LIMIT

| DEPTH(CM) | 27 CO 60 | 38 SR 90 | 55 CS 137 | 63 EU 155 | 94 PU 239 | 94 PU 240 | 94 PU 241 | 95 AM 241 | MASTER LOG |
|-----------|-----------|-----------|-----------|------------|-----------|-----------|-----------|------------|------------|
| 000-005 | 1.691E+00 | 1.380E+02 | 1.102E+02 | 2.341E+00 | 5.590E+00 | 6.185E+00 | 1.245E+02 | 6.131E+00 | 01032990 |
| 005-010 | 1.932E+00 | 2.343E+02 | 1.091E+02 | 2.818E+00 | 6.860E+00 | 7.590E+00 | 1.403E+02 | 9.671E+00 | 01033091 |
| 010-015 | 3.823E+00 | 3.183E+02 | 1.777E+02 | 6.059E+00 | 1.400E+01 | 1.514E+01 | 2.673E+02 | 1.874E+01 | 01033192 |
| 015-025 | 2.137E-01 | 2.304E+01 | 1.197E+01 | 1.6131E-02 | 2.422E-01 | 2.827E-01 | - | 2.601E-01 | 01033293 |
| 025-035 | 5.032E-01 | 1.240E+02 | 3.888E+01 | 1.783E-01 | 8.865E-01 | 1.087E+00 | 2.259E+01 | 9.811E-01 | 01033394 |
| 035-045 | 1.347E-01 | 1.231E+01 | 9.739E+00 | 1.4802E-02 | 1.281E-01 | 1.487E-01 | - | 1.958E-01 | 01033495 |
| 045-055 | 6.243E-02 | 2.493E+00 | 6.977E+00 | 1.455E-02 | 4.323E-02 | 4.955E-02 | 1.122E+00 | 1.2944E-02 | 01033596 |

501013

A-17

PROFILE TS0131

CONCENTRATIONS IN PCI/GM. MINUS FOR NO DATA. LESS THAN SIGN INDICATES INSTRUMENTAL DETECTION LIMIT

| DEPTH(CM) | 27 CO | 60 | 38 SR | 90 | 55 CS | 137 | 63 EU | 155 | 94 PU | 239 | 94 PU | 240 | 94 PU | 241 | 95 AM | 241 | MASTER LOG |
|-----------|------------|------------|-------|------------|-------|------------|-------|-----------|-------|-----------|-------|-----------|-------|------------|-------|----------|------------|
| 000-005 | 5.689E+00 | 3.421E+02 | | 3.404E+02 | | 7.559E+00 | | 1.806E+01 | | 2.033E+01 | | 3.914E+02 | | 2.383E+01 | | 01034190 | |
| 005-010 | 1.164E+00 | 1.689E+02 | | 6.374E+01 | | 1.081E+00 | | 2.488E+00 | | 2.886E+00 | | 5.275E+01 | | 3.433E+00 | | 01034291 | |
| 010-015 | 3.530E-01 | 1.279E+02 | | 1.269E+01 | | 1.073E-01 | | 4.047E-01 | | 4.892E-01 | | 8.950E+00 | | 4.577E-01 | | 01034392 | |
| 015-025 | ~3.644E-02 | 2.570E+01 | | 1.263E+00 | | ~3.121E-02 | | 4.860E-02 | | 6.153E-02 | | 1.272E+00 | | ~5.234E-02 | | 01034493 | |
| 025-035 | ~2.962E-02 | - | | 3.174E-01 | | ~1.845E-02 | | - | | - | | - | | ~3.118E-02 | | 01034594 | |
| 035-045 | ~3.832E-02 | 2.725E+00 | | 2.770E-01 | | ~2.029E-02 | | 3.036E-02 | | 3.586E-02 | | - | | ~3.728E-02 | | 01034695 | |
| 125-145 | ~1.573E-02 | ~3.604E+00 | | ~1.614E-02 | | ~1.365E-02 | | - | | - | | - | | ~2.359E-02 | | 01035101 | |

PROFILE TS0161

CONCENTRATIONS IN PCI/GM. MINUS FOR NO DATA, LESS THAN SIGN INDICATES INSTRUMENTAL DETECTION LIMIT

| DEPTH(CM) | 27 CO 60 | 38 SR 90 | 55 CS 137 | 63 EU 155 | 94 PU 239 | 94 PU 240 | 94 PU 241 | 95 AM 241 | MASTER LOG |
|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|------------|
| 000-005 | 4.068E-01 | 1.944E+01 | 5.658E+01 | 4.068E-01 | 1.114E+00 | 1.251E+00 | 2.686E+01 | 1.236E+00 | 01038290 |
| 005-010 | 5.977E-01 | 2.055E+01 | 3.322E+01 | 6.437E-01 | 1.442E+00 | 1.740E+00 | - | 1.655E+00 | 01038391 |
| 010-015 | 7.341E-01 | 3.382E+01 | 3.213E+01 | 7.821E-01 | 1.917E+00 | 2.118E+00 | 4.065E+01 | 2.209E+00 | 01038492 |
| 015-025 | 5.964E-01 | 3.122E+01 | 2.689E+01 | 6.284E-01 | 1.485E+00 | 1.675E+00 | 3.185E+01 | 1.641E+00 | 01038593 |
| 025-035 | 4.734E-01 | 4.104E+01 | 3.369E+01 | 5.829E-01 | 1.315E+00 | 1.482E+00 | 3.336E+01 | 1.498E+00 | 01038694 |
| 035-045 | 1.659E-01 | 2.481E+01 | 1.014E+01 | 1.391E-01 | 3.845E-01 | 4.383E-01 | 1.019E+01 | 4.236E-01 | 01038795 |
| 045-055 | 3.183E-01 | 3.909E+01 | 2.145E+01 | 5.077E-01 | 1.167E+00 | 1.273E+00 | 2.499E+01 | 1.112E+00 | 01038896 |
| 055-065 | 7.311E-02 | 2.249E+01 | 9.734E+00 | 4.180E-02 | 2.327E-01 | 2.412E-01 | - | 1.661E-01 | 01038997 |
| 065-075 | 2.418E-01 | 2.220E+01 | 1.492E+01 | 2.116E-01 | 6.077E-01 | 6.869E-01 | - | 5.977E-01 | 01039098 |
| 075-085 | 1.774E-01 | 1.888E+01 | 1.039E+01 | 1.616E-01 | 4.384E-01 | 4.784E-01 | 1.011E+01 | 4.946E-01 | 01039199 |
| 085-095 | 1.251E-01 | 2.479E+01 | 1.273E+01 | 3.405E-01 | 8.041E-01 | 8.928E-01 | - | 8.658E-01 | 01039201 |
| 095-105 | 4.389E-02 | 1.454E+01 | 7.014E+00 | 3.660E-02 | 1.622E-01 | 1.760E-01 | - | 6.667E-02 | 01039301 |
| 105-125 | 9.941E-01 | 4.429E+01 | 3.491E+01 | 1.295E+00 | 3.373E+00 | 3.727E+00 | 6.842E+01 | 3.258E+00 | 01039401 |
| 125-145 | 3.273E+00 | 1.808E+02 | 4.892E+01 | 4.644E+00 | 1.023E+01 | 1.087E+01 | 1.972E+02 | 1.178E+01 | 01039501 |
| 145-165 | 6.432E-02 | 1.095E+01 | 6.964E+00 | 3.911E-02 | 6.586E-02 | 8.216E-02 | 1.937E+00 | 7.113E-02 | 01039601 |
| 165-185 | 3.620E-02 | 3.153E-01 | 1.146E+00 | 2.180E-02 | - | - | - | 3.980E-02 | 01039701 |

PROFILE TS0171

CONCENTRATIONS IN PCI/GM. MINUS FOR NO DATA. LESS THAN SIGN INDICATES INSTRUMENTAL DETECTION LIMIT

| DEPTH(CM) | 27 CO 60 | 38 SR 90 | 55 CS 137 | 63 EU 155 | 94 PU 239 | 94 PU 240 | 94 PU 241 | 95 AM 241 | MASTER LOG |
|-----------|-----------|-----------|-----------|------------|-----------|-----------|-----------|------------|------------|
| 000-005 | 1.533E+00 | 1.360E+01 | 9.595E+01 | 2.206E+00 | 5.491E+00 | 6.212E+00 | 1.300E+02 | 6.667E+00 | 01027390 |
| 005-010 | 2.712E-01 | 5.063E+01 | 2.153E+01 | ~7.523E-02 | 4.748E-01 | 5.986E-01 | - | 5.423E-01 | 01027491 |
| 010-015 | 1.953E-01 | 4.138E+01 | 1.719E+01 | ~6.581E-02 | 3.323E-01 | 4.413E-01 | 8.955E+00 | 3.982E-01 | 01027592 |
| 015-025 | 1.796E-01 | 3.558E+01 | 4.095E+00 | ~5.441E-02 | - | - | - | 4.563E-01 | 01027693 |
| 025-035 | 7.023E-02 | 3.690E+00 | 1.407E+00 | ~3.935E-02 | - | - | - | ~6.851E-02 | 01027794 |
| 035-045 | 6.275E-02 | 3.438E+00 | 1.118E+00 | ~1.987E-02 | - | - | - | 1.592E-01 | 01027895 |

PROFILE TS0181

CONCENTRATIONS IN PCI/GM. MINUS FOR NO DATA. LESS THAN SIGN INDICATES INSTRUMENTAL DETECTION LIMIT

| DEPTH(CM) | 27 CO 60 | 38 SR 90 | 55 CS 137 | 63 EU 155 | 94 PU 239 | 94 PU 240 | 94 PU 241 | 95 AM 241 | MASTER LOG |
|-----------|------------|-----------|-----------|------------|-----------|-----------|-----------|------------|------------|
| 000-005 | 1.939E+00 | 1.202E+02 | 2.838E+01 | 2.257E+00 | 6.050E+00 | 6.640E+00 | 1.350E+02 | 6.010E+00 | 01062690 |
| 005-010 | 1.001E+00 | 9.784E+01 | 2.225E+01 | 9.557E-01 | 2.481E+00 | 2.800E+00 | 4.946E+01 | 2.605E+00 | 01062791 |
| 010-015 | 2.397E+00 | 1.732E+02 | 3.185E+01 | 2.836E+00 | 6.842E+00 | 7.617E+00 | 1.385E+02 | 7.218E+00 | 01062892 |
| 015-025 | 5.609E-01 | 6.063E+01 | 5.205E+00 | 1.433E-01 | 8.968E-01 | 1.134E+00 | - | 9.761E-01 | 01062993 |
| 025-035 | 1.479E-01 | 4.141E+01 | 1.398E+00 | ~3.397E-02 | 1.641E-01 | 2.128E-01 | 4.117E+00 | 1.614E-01 | 01063094 |
| 035-045 | 1.068E-01 | 3.374E+01 | 1.266E+00 | ~3.360E-02 | 1.545E-01 | 2.051E-01 | 3.832E+00 | ~6.023E-02 | 01063195 |
| 045-055 | 1.560E-01 | 6.252E+01 | 1.517E+01 | 5.523E-02 | 1.734E-01 | 2.326E-01 | - | 1.739E-01 | 01063296 |
| 055-065 | ~4.328E-02 | 2.918E+01 | 1.005E+01 | ~4.523E-02 | 7.932E-02 | 1.025E-01 | 1.910E+00 | ~7.878E-02 | 01063397 |
| 065-085 | ~5.306E-02 | 3.470E+00 | 1.503E+00 | ~2.426E-02 | - | - | - | ~4.505E-02 | 01063401 |

५८

| PROFILE TS0191 | | | | | | | | | | | | | | | | | | | | | | | | | |
|--|---------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|----------|----|-----|----|----|-----|----|----|-----|----|----|-----|----|----|-----|------------|
| CONCENTRATIONS IN PCI/GM. MINUS FOR NO DATA. LESS THAN SIGN INDICATES INSTRUMENTAL DETECTION LIMIT | | | | | | | | | | | | | | | | | | | | | | | | | |
| DEPTH(CM) | 27 | CO | 60 | 38 | SR | 90 | 55 | CS | 137 | 63 | EU | 155 | 94 | PU | 239 | 94 | PU | 240 | 94 | PU | 241 | 95 | AM | 241 | MASTER LOG |
| A-20 | 000-005 | 8.743E-01 | 4.950E+01 | 6.207E+01 | 1.181E+00 | 2.682E+00 | 3.135E+00 | 3.365E+01 | 3.109E+00 | 01060090 | | | | | | | | | | | | | | | |
| | 005-010 | 6.950E-01 | 5.117E+01 | 3.766E+01 | 1.044E+00 | 2.262E+00 | 2.529E+00 | 4.451E+01 | 2.420E+00 | 01060191 | | | | | | | | | | | | | | | |
| | 010-015 | 6.680E-01 | 4.806E+01 | 3.782E+01 | 8.104E-01 | 1.492E+00 | 1.697E+00 | 3.727E+01 | 2.013E+00 | 01060292 | | | | | | | | | | | | | | | |
| | 015-025 | 1.404E-01 | 2.204E+01 | 8.203E+00 | 1.249E-01 | 3.218E-01 | 4.127E-01 | - | 3.175E-01 | 01060393 | | | | | | | | | | | | | | | |
| | 025-035 | 2.829E-01 | 2.595E+01 | 2.228E+01 | 3.545E-01 | 8.635E-01 | 9.617E-01 | 1.800E+01 | 8.644E-01 | 01060494 | | | | | | | | | | | | | | | |
| | 035-045 | 1.614E-01 | 1.119E+01 | 5.955E+00 | 1.973E-01 | 4.388E-01 | 4.986E-01 | - | 4.842E-01 | 01060595 | | | | | | | | | | | | | | | |
| | 045-055 | 4.577E-01 | 4.060E+01 | 2.106E+01 | 4.901E-01 | 1.285E+00 | 1.475E+00 | - | 1.323E+00 | 01060695 | | | | | | | | | | | | | | | |
| | 055-065 | 8.595E-01 | 6.559E+01 | 4.649E+01 | 7.793E-01 | 2.006E+00 | 2.252E+00 | 4.323E+01 | 2.183E+00 | 01060797 | | | | | | | | | | | | | | | |
| | 065-075 | 1.712E+00 | 1.155E+02 | 5.023E+01 | 2.156E+00 | 4.946E+00 | 5.428E+00 | - | 4.869E+00 | 01060898 | | | | | | | | | | | | | | | |
| | 075-085 | 8.392E-01 | 8.829E+01 | 2.723E+01 | 6.964E-01 | 2.051E+00 | 2.380E+00 | 4.423E+01 | 2.159E+00 | 01060999 | | | | | | | | | | | | | | | |
| | 085-095 | 1.254E-01 | 3.598E+01 | 2.389E+01 | 1.231E-01 | 3.367E-01 | 3.762E-01 | 6.167E+00 | 4.027E-01 | 01061001 | | | | | | | | | | | | | | | |
| | 095-105 | 5.140E-02 | 9.279E+00 | 1.433E+01 | 5.739E-02 | 1.872E-01 | 2.080E-01 | 3.787E+00 | 2.439E-01 | 01061101 | | | | | | | | | | | | | | | |
| | 105-115 | 1.687E-01 | 1.499E+01 | 2.808E+01 | 1.776E-01 | 4.467E-01 | 5.207E-01 | - | 3.863E-01 | 01061201 | | | | | | | | | | | | | | | |
| | 115-125 | 2.271E-02 | 2.384E+00 | 3.481E+00 | 1.979E-02 | 4.568E-03 | 5.766E-03 | - | 3.873E-02 | 01061301 | | | | | | | | | | | | | | | |

PROFILE TS0201

CONCENTRATIONS IN PC1/GM. MINUS FOR NO DATA. LESS THAN SIGN INDICATES INSTRUMENTAL DETECTION LIMIT

| DEPTH(CM) | 27 CO 60 | 38 SR 90 | 55 CS 137 | 63 EU 155 | 94 PU 239 | 94 PU 240 | 94 PU 241 | 95 AM 241 | MASTER LOG |
|-----------|------------|-----------|-----------|------------|-----------|-----------|-----------|------------|------------|
| 000-005 | 9.937E-01 | 3.451E+01 | 5.027E+01 | 1.241E+00 | 2.926E+00 | 3.515E+00 | 7.482E+01 | 3.645E+00 | 01061590 |
| 005-010 | 1.149E+00 | 4.500E+01 | 2.882E+01 | 1.275E+00 | 3.740E+00 | 4.563E+00 | 8.973E+01 | 4.022E+00 | 01061691 |
| 010-015 | 4.761E-01 | 2.486E+01 | 1.514E+01 | 3.756E-01 | 9.673E-01 | 1.155E+00 | 2.188E+01 | 1.126E+00 | 01061792 |
| 015-025 | 5.455E-02 | 2.057E+01 | 1.002E+01 | ~3.528E-02 | - | - | - | ~6.568E-02 | 01061893 |
| 025-035 | ~5.225E-02 | 1.723E+01 | 7.171E+00 | ~3.464E-02 | - | - | - | ~7.027E-02 | 01061994 |
| 035-045 | ~2.385E-02 | - | 1.063E+00 | ~2.079E-02 | - | - | - | ~3.775E-02 | 01062095 |
| 045-055 | ~2.560E-02 | - | 5.964E-01 | ~1.922E-02 | - | - | - | ~3.550E-02 | 01062196 |

PROFILE TS0231

CONCENTRATIONS IN PC1/GM. MINUS FOR NO DATA. LESS THAN SIGN INDICATES INSTRUMENTAL DETECTION LIMIT

| DEPTH(CM) | 27 CO 60 | 38 SR 90 | 55 CS 137 | 63 EU 155 | 94 PU 239 | 94 PU 240 | 94 PU 241 | 95 AM 241 | MASTER LOG |
|-----------|------------|-----------|-----------|------------|-----------|-----------|-----------|------------|------------|
| 000-005 | 5.856E-01 | 2.997E+01 | 3.506E+01 | 7.239E-01 | 1.700E+00 | 1.896E+00 | 4.806E+01 | 1.957E+00 | 01036790 |
| 005-010 | 5.329E-02 | 1.078E+01 | 1.651E+01 | ~4.887E-02 | 2.152E-01 | 2.436E-01 | 5.856E+00 | ~8.464E-02 | 01036891 |
| 010-015 | ~3.532E-02 | 9.788E+00 | 4.106E+01 | ~6.437E-02 | 1.377E-01 | 1.532E-01 | - | ~1.167E-01 | 01036992 |
| 015-025 | ~3.628E-02 | 1.275E+01 | 4.408E+00 | ~3.018E-02 | 1.475E+00 | 1.653E+00 | 3.179E+01 | ~5.923E-02 | 01037093 |
| 025-035 | ~4.251E-02 | 1.088E+01 | 1.886E+00 | ~3.301E-02 | 2.904E-02 | 3.756E-02 | - | ~5.369E-02 | 01037194 |
| 035-045 | ~2.038E-02 | 3.343E+00 | 1.454E+00 | ~4.443E-02 | 8.559E-03 | 1.040E-02 | - | ~6.500E-02 | 01037295 |

5010110

A-22

PROFILE TS0241

CONCENTRATIONS IN PC1/GM. MINUS FOR NO DATA. LESS THAN SIGN INDICATES INSTRUMENTAL DETECTION LIMIT

| DEPTH(CM) | 27 | CO | 60 | 38 | SR | 90 | 55 | CS | 137 | 63 | EU | 155 | 94 | PU | 239 | 94 | PU | 240 | 94 | PU | 241 | 95 | AM | 241 | MASTER LOG |
|-----------|-----------|----|----|-----------|----|----|-----------|----|-----------|----|-----------|-----|-----------|----|-----------|----|-----------|-----|----------|----|-----|----|----|-----|------------|
| 000-005 | 6.473E-02 | | | 3.286E+00 | | | 6.068E+00 | | 1.444E-01 | | 1.826E+00 | | 2.212E+00 | | 3.336E+01 | | 2.109E-01 | | 06067890 | | | | | | |
| 005-010 | 6.014E-02 | | | 3.555E+00 | | | 2.915E+00 | | 1.394E-01 | | 2.663E-01 | | 2.789E-01 | | 5.622E+00 | | 2.445E-01 | | 06067991 | | | | | | |
| 010-015 | 8.766E-02 | | | 4.617E+00 | | | 2.336E+00 | | 1.518E-01 | | 3.093E-01 | | 3.371E-01 | | 7.775E+00 | | 2.302E-01 | | 06068092 | | | | | | |
| 015-025 | 9.320E-02 | | | 3.643E+00 | | | 3.020E+00 | | 1.667E-01 | | 2.982E-01 | | 3.159E-01 | | - | | 3.940E-01 | | 06068193 | | | | | | |
| 025-035 | 1.498E-01 | | | 5.432E+00 | | | 2.487E+00 | | 2.574E-01 | | 3.571E-01 | | 3.917E-01 | | 8.090E+00 | | 5.050E-01 | | 06068294 | | | | | | |
| 035-045 | 1.157E-01 | | | 7.428E+00 | | | 2.229E+00 | | 1.952E-01 | | 5.545E-01 | | 5.955E-01 | | 9.608E+00 | | 4.475E-01 | | 06068395 | | | | | | |

PROFILE TS0251

CONCENTRATIONS IN PC1/GM. MINUS FOR NO DATA. LESS THAN SIGN INDICATES INSTRUMENTAL DETECTION LIMIT

| DEPTH(CM) | 27 | CO | 60 | 38 | SR | 90 | 55 | CS | 137 | 63 | EU | 155 | 94 | PU | 239 | 94 | PU | 240 | 94 | PU | 241 | 95 | AM | 241 | MASTER LOG |
|-----------|-----------|----|----|-----------|----|----|-----------|----|-----------|----|-----------|-----|-----------|----|-----------|----|-----------|-----|----------|----|-----|----|----|-----|------------|
| 000-005 | 1.780E-01 | | | 8.779E+00 | | | 1.258E+01 | | 1.778E-01 | | 3.723E-03 | | 4.977E-03 | | - | | 4.207E-01 | | 06066490 | | | | | | |
| 005-010 | 1.893E-01 | | | 7.572E+00 | | | 7.356E+00 | | 2.686E-01 | | 4.577E-01 | | 5.086E-01 | | - | | 4.323E-01 | | 06066591 | | | | | | |
| 010-015 | 2.064E-01 | | | 8.239E+00 | | | 6.185E+00 | | 4.306E-01 | | 5.140E-01 | | 5.577E-01 | | 1.019E+01 | | 8.505E-01 | | 06066692 | | | | | | |
| 015-025 | 1.609E-01 | | | 6.207E+00 | | | 7.320E+00 | | 2.743E-01 | | 9.198E-01 | | 1.048E+00 | | - | | 6.198E-01 | | 06066793 | | | | | | |
| 025-035 | 2.093E-01 | | | 1.104E+01 | | | 3.476E+00 | | 4.122E-01 | | 6.662E-01 | | 5.500E-01 | | - | | 9.329E-01 | | 06066894 | | | | | | |
| 035-045 | 3.101E-01 | | | 1.797E+01 | | | 4.063E+00 | | 6.306E-01 | | 1.124E+00 | | 1.262E+00 | | 2.529E+01 | | 1.451E+00 | | 06066995 | | | | | | |

PROFILE TS0261

CONCENTRATIONS IN PCI/GM. MINUS FOR NO DATA. LESS THAN SIGN INDICATES INSTRUMENTAL DETECTION LIMIT

| DEPTH(CM) | 27 CO 60 | 38 SR 90 | 55 CS 137 | 63 EU 155 | 94 PU 239 | 94 PU 240 | 94 PU 241 | 95 AM 241 | MASTER LOG |
|-----------|-----------|-----------|-----------|-------------|-----------|-----------|-----------|-----------|------------|
| 000-005 | 1.595E-01 | 4.689E+00 | 7.892E+00 | 1.339E-01 | 4.147E-01 | 4.775E-01 | - | 4.250E-01 | 06065290 |
| 005-010 | 1.714E-01 | 6.419E+00 | 3.718E+00 | 2.119E-01 | 4.304E-01 | 5.108E-01 | 1.099E+01 | 4.833E-01 | 06065391 |
| 010-015 | 1.820E-01 | 7.320E+00 | 3.703E+00 | 1.273E-01 | 5.171E-01 | 6.045E-01 | - | 4.396E-01 | 06065492 |
| 015-025 | 1.195E-01 | 1.137E+01 | 4.421E+00 | 2.5.077E-02 | 3.477E-01 | 4.014E-01 | - | 3.377E-01 | 06065593 |
| 025-035 | 7.167E-02 | 8.608E+00 | 1.776E+00 | 2.4.514E-02 | 5.689E-02 | 6.640E-02 | - | 2.351E-02 | 06065694 |
| 035-045 | 2.022E-02 | 4.395E+00 | 1.572E-01 | 2.3.395E-02 | - | - | - | 2.234E-02 | 06065795 |
| 045-055 | 2.555E-02 | 2.903E-01 | 2.413E-02 | 2.1.532E-02 | - | - | - | 2.670E-02 | 06065896 |
| 055-065 | 1.574E-02 | 3.916E-01 | 2.870E-02 | 2.1.470E-02 | 2.965E-03 | 3.395E-03 | - | 2.626E-02 | 06065901 |
| 065-085 | 2.278E-02 | 3.153E-01 | 2.787E-02 | 2.2.003E-02 | - | - | - | 3.491E-02 | 06066001 |
| 085-105 | 1.685E-02 | 4.505E-01 | 2.304E-02 | 2.1.706E-02 | - | - | - | 2.113E-02 | 06066101 |
| 105-125 | 1.386E-02 | 2.703E-01 | 2.129E-02 | 2.1.441E-02 | - | - | - | 2.522E-02 | 06066201 |
| 125-145 | 2.914E-02 | 2.928E-01 | 2.610E-02 | 2.1.732E-02 | - | - | - | 3.130E-02 | 06066301 |

PROFILE TS0271

CONCENTRATIONS IN PCI/GM. MINUS FOR NO DATA. LESS THAN SIGN INDICATES INSTRUMENTAL DETECTION LIMIT

| DEPTH(CM) | 27 CO 60 | 38 SR 90 | 55 CS 137 | 63 EU 155 | 94 PU 239 | 94 PU 240 | 94 PU 241 | 95 AM 241 | MASTER LOG |
|-----------|-----------|-----------|-----------|-------------|-----------|-----------|-----------|-----------|------------|
| 000-005 | 1.185E-01 | 2.428E+00 | 4.257E+00 | 9.541E-02 | 3.604E-01 | 4.093E-01 | 7.315E+00 | 2.480E-01 | 06089190 |
| 005-010 | 3.403E-02 | 1.789E+00 | 1.605E+00 | 1.008E-01 | 3.768E-01 | 4.354E-01 | 8.505E+00 | 2.150E-01 | 06089291 |
| 010-015 | 1.109E-01 | 1.533E+00 | 1.121E+00 | 1.192E-01 | 4.159E-01 | 4.711E-01 | 9.383E+00 | 2.951E-01 | 06089392 |
| 015-025 | 3.383E-02 | 4.123E-01 | 2.725E-01 | 2.5.347E-02 | - | - | - | 2.086E-02 | 06089493 |
| 025-035 | 5.716E-02 | 3.153E-01 | 1.800E-01 | 2.5.288E-02 | - | - | - | 2.261E-02 | 06089594 |
| 035-045 | 3.791E-02 | 3.604E-01 | 1.194E-01 | 2.5.671E-02 | 6.563E-03 | 6.946E-03 | - | 2.766E-02 | 06089695 |

5010142-

PROFILE TS0301

CONCENTRATIONS IN PCI/GM. MINUS FOR NO DATA. LESS THAN SIGN INDICATES INSTRUMENTAL DETECTION LIMIT


| DEPTH(CM) | 27 CO | 60 | 38 SR | 90 | 55 CS | 137 | 63 EU | 155 | 94 PU | 239 | 94 PU | 240 | 94 PU | 241 | 95 AM | 241 | MASTER LOG |
|-----------|---------|-----------|-------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|------------|
| A-24 | 000-005 | 2.592E-01 | | 1.776E+01 | 1.136E+01 | | 5.649E-01 | | 1.344E+00 | | 1.444E+00 | | 2.413E+01 | | 1.023E+00 | | 06093290 |
| | 005-010 | 3.294E-01 | | 1.574E+01 | | 9.599E+00 | | 4.761E-01 | | 1.102E+00 | | 1.218E+00 | | - | | 9.770E-01 | 06093391 |
| | 010-015 | 3.338E-01 | | 1.489E+01 | | 9.770E+00 | | 5.149E-01 | | 1.093E+00 | | 1.160E+00 | | 2.315E+01 | | 1.036E+00 | 06093492 |
| | 015-025 | 2.459E-01 | | 1.611E+01 | | 1.055E+01 | | 4.775E-01 | | 1.023E+00 | | 1.111E+00 | | 2.023E+01 | | 1.155E+00 | 06093593 |
| | 025-035 | 5.541E-01 | | 3.409E+01 | | 1.778E+01 | | 1.276E+00 | | 2.888E+00 | | 3.083E+00 | | 5.658E+01 | | 3.002E+00 | 06093694 |
| | 035-045 | 3.004E-01 | | 2.146E+01 | | 9.856E+00 | | 6.081E-01 | | 1.244E+00 | | 1.350E+00 | | - | | 9.239E-01 | 06093795 |





Appendix B. The Distribution of Radionuclides with Depth in Soil Profiles of Bikini and Eneu Islands


The following figures graphically present the concentrations of five selected radionuclides (^{60}Co , ^{90}Sr , ^{137}Cs , ^{239}Pu , and ^{241}Am) with depth of soil. One graph is given for each sample location of the islands (see Figs. 2, 3) and each corresponds to the tabular presentation of the same data in Appendix A. Throughout, open symbols indicate detection limits and solid symbols indicate measured values: $\blacktriangledown = ^{60}\text{Co}$, $\blacktriangle = ^{90}\text{Sr}$, $\bullet = ^{137}\text{Cs}$, $\blacksquare = ^{239}\text{Pu}$, and $\blacklozenge = ^{241}\text{Am}$. Figures are grouped according to general location of the samples.




5010144





B-2



























5010151



B-10





Location





5010152







B-11

5010153







Appendix C. Geographical Distribution of Radioactivity in the Surface Soil (0 to 15 cm) of Bikini and Eneu Islands

The following maps present the concentrations of ^{60}Co , ^{90}Sr , ^{137}Cs , ^{239}Pu , ^{240}Pu , and ^{241}Am (pCi/g dry weight) at the various samplings sites of the top 15 cm of soil on Bikini and Eneu Islands.


5010157

BLANK PAGE

-3 5010158



5010159



in Island, 60^{Co}

۳۱


5010100





-J U T U + U ^ 2










5010166







Bikini Isl.
C-8




5010170

E_b




5010171

land, 60
Co
C-9



5010172



C-1

5010173



5010174


G-11

and, 137_{CS}



1

5010175



501017b




Enneu Island


८

5010177

t_{pu}



5010178



5010179



5010180



Lneu Island

C-

5010181

5010182



EVALUATION OF THE RADIOLOGICAL QUALITY OF THE WATER ON BIKINI AND ENEU ISLANDS IN 1975: DOSE ASSESSMENT BASED ON INITIAL SAMPLING


V. E. Noshkin, W. L. Robison,
K. M. Wong, and R. J. Eagle

January 21, 1977

Prepared for U.S. Energy Research & Development
Administration under contract No. W-7405-Eng-48



**LAWRENCE
LIVERMORE
LABORATORY**
University of California / Livermore



NOTICE

This report was prepared as an account of work sponsored by the United States Government. Neither the United States nor the United States Energy Research & Development Administration, nor any of their employees, nor any of their contractors, subcontractors, or their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness or usefulness of any information, apparatus, product or process disclosed, or represents that its use would not infringe privately-owned rights

NOTICE

Reference to a company or product name does not imply approval or recommendation of the product by the University of California or the U.S. Energy Research & Development Administration to the exclusion of others that may be suitable

Printed in the United States of America

Available from

National Technical Information Service
U S Department of Commerce
5285 Port Royal Road
Springfield, VA 22161

Price Printed Copy \$, Microfiche \$3.00

| Page Range | Domestic Price | Page Range | Domestic Price |
|------------|----------------|------------|----------------|
| 001-025 | \$ 3.50 | 326-350 | 10.00 |
| 026-050 | 4.00 | 351-375 | 10.50 |
| 051-075 | 4.50 | 376-400 | 10.75 |
| 076-100 | 5.00 | 401-425 | 11.00 |
| 101-125 | 5.50 | 426-450 | 11.75 |
| 126-150 | 6.00 | 451-475 | 12.00 |
| 151-175 | 6.75 | 476-500 | 12.50 |
| 176-200 | 7.50 | 501-525 | 12.75 |
| 201-225 | 7.75 | 526-550 | 13.00 |
| 226-250 | 8.00 | 551-575 | 13.50 |
| 251-275 | 9.00 | 576-600 | 13.75 |
| 276-300 | 9.25 | 601-up | * |
| 301-325 | 9.75 | | |

*Add \$2.50 for each additional 100 page increment from 601 to 1,000 pages;
add \$4.50 for each additional 100 page increment over 1,000 pages

5010164